Fredrik Dahlqvist \square

University College London, United Kingdom

Renato Neves \square

University of Minho & INESC-TEC, Portugal

— Abstract

Programs with a *continuous* state space or that interact with physical processes often require notions of equivalence going beyond the standard binary setting in which equivalence either holds or does not hold. In this paper we explore the idea of equivalence taking values in a quantale \mathcal{V} , which covers the cases of (in)equations and (ultra)metric equations among others.

Our main result is the introduction of a \mathcal{V} -equational deductive system for linear λ -calculus together with a proof that it is sound and complete (in fact, an internal language) for a class of enriched autonomous categories. In the case of inequations, we get an internal language for autonomous categories enriched over partial orders. In the case of (ultra)metric equations, we get an internal language for autonomous categories enriched over (ultra)metric spaces.

We use our results to obtain examples of inequational and metric equational systems for higherorder programs that contain real-time and probabilistic behaviour.

2012 ACM Subject Classification Theory of computation \rightarrow Models of computation

Keywords and phrases λ -calculus, enriched category theory, quantale, equational theory

Acknowledgements This work was financed by the ERDF – European Regional Development Fund through the Operational Programme for Competitiveness and Internationalisation - COMPETE 2020 Programme and by National Funds through the Portuguese funding agency, FCT - Fundação para a Ciência e a Tecnologia, within project POCI-01-0145-FEDER-030947. This work was also financed by the UK Research Institute in Verified Trustworthy Software Systems, via project "Quantitative Algebraic Reasoning for Hybrid Programs", and by the Leverhulme Project Grant "Verification of Machine Learning Algorithms". The authors are very grateful for the reviewer's incisive feedback.

1 Introduction

Programs frequently act over a *continuous* state space or interact with physical processes like time progression or the movement of a vehicle. Such features naturally call for notions of approximation and refinement integrated in different aspects of program equivalence. Our paper falls in this line of research. Specifically, our aim is to integrate notions of approximation and refinement into the *equational system* of linear λ -calculus [4, 27, 28].

The core idea that we explore in this paper is to have equations $t =_q s$ labelled by elements q of a quantale \mathcal{V} . This covers a wide range of situations, among which the cases of (in)equations [22, 2] and metric equations [29, 30]. The latter case is perhaps less known: it consists of equations $t =_{\epsilon} s$ labelled by a non-negative rational number ϵ which represents the 'maximum distance' that the two terms t and s can be from each other. In order to illustrate metric equations, consider a programming language with a (ground) type X and a signature of operations $\Sigma = \{ \texttt{wait}_n : X \to X \mid n \in \mathbb{N} \}$ that model time progression over computations of type X. Specifically, $\texttt{wait}_n(x)$ reads as "add a latency of n seconds to the computation x". In this context, the following axioms involving metric equations arise naturally:

$$wait_0(x) =_0 x \qquad wait_n(wait_m(x)) =_0 wait_{n+m}(x) \qquad \frac{\epsilon = |m-n|}{wait_n(x) =_{\epsilon} wait_m(x)}$$
(1)

An equation $t =_0 s$ states that the terms t and s are exactly the same and equations $t =_{\epsilon} s$ state that t and s differ by at most ϵ seconds in their execution time.

Contributions. In this paper we introduce an equational deductive system for linear λ calculus in which equations are labelled by elements of a quantale \mathcal{V} . By using key features of
a quantale's structure, we show that this deductive system is *sound and complete* for a class
of enriched symmetric monoidal closed categories (*i.e.* enriched autonomous categories). In
particular, if we fix \mathcal{V} to be the Boolean quantale this class of categories consists of autonomous
categories enriched over partial orders. If we fix \mathcal{V} to be the (ultra)metric quantale, this
class of categories consists of autonomous categories enriched over (ultra)metric spaces. The
aforementioned example of wait calls fits in the setting in which \mathcal{V} is the metric quantale.
Our result provides this example with a sound and complete metric equational system, where
the models are all those autonomous categories enriched over metric spaces that can soundly
interpret the axioms of wait calls (1).

The next contribution of our paper falls in one of the major topics of categorical logic: to establish *logical descriptions* of certain classes of categories. A famous result of this kind is the correspondence between λ -calculus and Cartesian closed categories which states that the former is the *internal language* of the latter [23] – such a correspondence allows to study Cartesian closed categories by means of logical tools. An analogous result is presented in [27, 28] for linear λ -calculus and symmetric monoidal closed (*i.e.* autonomous) categories. We show that linear λ -calculus equipped with a \mathcal{V} -equational system is the internal language of autonomous categories enriched over 'generalised metric spaces'.

Outline. Section 2 recalls linear λ -calculus, its equational system, and the famous correspondence to autonomous categories, via soundness, completeness, and internal language theorems. The contents of this section are slight adaptations of results presented in [27, 4], the main difference being that we forbid the exchange rule to be explicitly part of linear λ -calculus (instead it is only admissible). This choice is important to ensure that judgements in the calculus have *unique* derivations, which allows us to refer to their interpretations unambiguously [37]. Section 3 presents the main contributions of this paper. It walks a path analogous to Section 2, but now in the setting of \mathcal{V} -equations (*i.e.* equations labelled by elements of a quantale \mathcal{V}). As we will see, the semantic counterpart of moving from equations to \mathcal{V} -equations is to move from categories to categories enriched over \mathcal{V} -categories. The latter, often regarded as generalised metric spaces, are central entities in a fruitful area of enriched category theory that aims to treat uniformly different kinds of 'structured sets', such as partial orders, fuzzy partial orders, and (ultra)metric spaces [24, 38, 39]. Our results are applicable to all these examples. Section 4 presents some examples of \mathcal{V} -equational axioms and corresponding models. Specifically, we will revisit the axioms of wait calls (1) and consider an inequational variant. Then we will study a metric axiom for probabilistic programs and show that the category of Banach spaces and short linear maps is a model for the resulting metric theory. We will additionally use this example to illustrate how our deductive system allows to compute an approximate distance between two probabilistic programs easily as opposed to computing an exact distance 'semantically' which tends to involve quite complex operators. Finally, Section 5 establishes a functorial connection between our results and previous well-known semantics for linear logic [10, 28], and concludes with a brief exposition of future work. We assume knowledge of λ -calculus and category theory [27, 28, 23, 26]. Proofs omitted in the main text are available in the appendix.

Related work. Several approaches to incorporating quantitative information to programming languages have been explored in the literature. Closest to this work are various approaches targeted at λ -calculi. In [7, 8] a notion of distance called *context distance* is

developed, first for an affine, then for a more general λ -calculus, with probabilistic programs as the main motivation. [13] considers a notion of quantale-valued *applicative (bi)similarity*, an operational *coinductive* technique used for showing contextual equivalence between two programs. Recently, [33] presented several Cartesian closed categories of generalised metric spaces that provide a quantitative semantics to simply-typed λ -calculus based on a generalisation of logical relations. None of these examples reason about distances in a quantitative equational system, and in this respect our work is closer to the metric universal algebra developed in [29, 30].

A different approach consists in encoding quantitative information via a type system. In particular, graded (modal) types [15, 12, 31] have found applications in *e.g.* differential privacy [34] and information flow [1]. This approach is to some extent orthogonal to ours as it mainly aims to model coeffects, whilst we aim to reason about the intrinsic quantitative nature of λ -terms acting *e.g.* on continuous or ordered spaces.

Quantum programs provide an interesting example of intrinsically quantitative programs, by which we mean that the metric structure on quantum states does not arise from (co)effects. Recently, [18] showed how the issue of noise in a quantum while-language can be handled by developing a *deductive system* to determine how similar a quantum program is from its idealised, noise-free version; an approach very much in the spirit of this work.

2 An internal language for autonomous categories

In this section we briefly recall linear λ -calculus, which can be regarded as a term assignment system for the exponential free, multiplicative fragment of *intuitionistic linear logic*. Then we recall that it is sound and complete w.r.t. autonomous categories, and also that it is an internal language for such categories. We mention only what is needed to present our results, the interested reader will find a more detailed exposition in [27, 4, 28]. Let us start by fixing a *class G* of ground types. The grammar of types for linear λ -calculus is given by:

$$\mathbb{A} ::= X \in G \mid \mathbb{I} \mid \mathbb{A} \otimes \mathbb{A} \mid \mathbb{A} \multimap \mathbb{A}$$

We also fix a class Σ of sorted operation symbols $f : \mathbb{A}_1, \ldots, \mathbb{A}_n \to \mathbb{A}$ with $n \geq 1$. As usual, we use Greek letters $\Gamma, \Delta, E, \ldots$ to denote *typing contexts*, *i.e.* lists $x_1 : \mathbb{A}_1, \ldots, x_n : \mathbb{A}_n$ of typed variables such that each variable x_i occurs at most once in x_1, \ldots, x_n .

We will use the notion of a shuffle for building a linear typing system such that the exchange rule is admissible and each judgement $\Gamma \triangleright v : \mathbb{A}$ (details about these below) has a unique derivation – this will allow us to refer to a judgement's denotation $[\Gamma \triangleright v : \mathbb{A}]$ unambiguously. By shuffle we mean a permutation of typed variables in a context sequence $\Gamma_1, \ldots, \Gamma_n$ such that for all $i \leq n$ the relative order of the variables in Γ_i is preserved [37]. For example, if $\Gamma_1 = x : \mathbb{A}, y : \mathbb{B}$ and $\Gamma_2 = z : \mathbb{C}$ then $z : \mathbb{C}, x : \mathbb{A}, y : \mathbb{B}$ is a shuffle but $y : \mathbb{B}, x : \mathbb{A}, z : \mathbb{C}$ is not, because we changed the order in which x and y appear in Γ_1 . As explained in [37] (and also in the proof of Lemma 1), such a restriction on relative orders is crucial for judgements having unique derivations. We denote by $\mathrm{Sf}(\Gamma_1; \ldots; \Gamma_n)$ the set of shuffles on $\Gamma_1, \ldots, \Gamma_n$.

The term formation rules of linear λ -calculus are listed in Fig. 1. They correspond to the natural deduction rules of the exponential free, multiplicative fragment of intuitionistic linear logic.

▶ Lemma 1. All judgements $\Gamma \triangleright v : \mathbb{A}$ have a unique derivation.

Substitution is defined in the expected way, and the following result is standard.

▶ Lemma 2 (Exchange and Substitution). For every judgement $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \triangleright v : \mathbb{C}$ we can derive $\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \triangleright v : \mathbb{C}$. For all judgements $\Gamma, x : \mathbb{A} \triangleright v : \mathbb{B}$ and $\Delta \triangleright w : \mathbb{A}$ we can derive $\Gamma, \Delta \triangleright v[w/x] : \mathbb{B}$.

We now recall the interpretation of judgements $\Gamma \triangleright v : \mathbb{A}$ in a symmetric monoidal closed (autonomous) category C . But before proceeding with this description, let us fix notation for some of the constructions available in autonomous categories. For all C -objects X, Y, Z, $\mathsf{sw} : X \otimes Y \to Y \otimes X$ denotes the symmetry morphism, $\lambda : \mathbb{I} \otimes X \to X$ the left unitor, $\operatorname{app} : (X \multimap Y) \otimes X \to Y$ the application, and $\alpha : X \otimes (Y \otimes Z) \to (X \otimes Y) \otimes Z$ the left associator. Moreover for all C -morphisms $f : X \otimes Y \to Z$ we denote the corresponding curried version (right transpose) by $\overline{f} : X \to (Y \multimap Z)$.

For all ground types $X \in G$ we postulate an interpretation $\llbracket X \rrbracket$ as a C-object. Types are then interpreted by induction over the type structure of linear λ -calculus, using the tensor \otimes and exponential \multimap constructs of autonomous categories. Given a non-empty context $\Gamma = \Gamma', x : \mathbb{A}$, its interpretation is defined by $\llbracket \Gamma', x : \mathbb{A} \rrbracket = \llbracket \Gamma' \rrbracket \otimes \llbracket \mathbb{A} \rrbracket$ if Γ' is non-empty and $\llbracket \Gamma', x : \mathbb{A} \rrbracket = \llbracket \mathbb{A} \rrbracket$ otherwise. The empty context - is interpreted as $\llbracket - \rrbracket = \rrbracket$ where \rrbracket is the unit of \otimes in C. To keep notation simple, given $X_1, \ldots, X_n \in \mathsf{C}$ we write $X_1 \otimes \cdots \otimes X_n$ for the *n*-tensor $(\ldots (X_1 \otimes X_2) \otimes \ldots) \otimes X_n$, and similarly for C-morphisms.

We will also need some 'housekeeping' morphisms to handle interactions between context interpretation and the autonomous structure of **C**. Specifically, given contexts $\Gamma_1, \ldots, \Gamma_n$ we denote by $\operatorname{sp}_{\Gamma_1;\ldots;\Gamma_n} : \llbracket\Gamma_1, \ldots, \Gamma_n \rrbracket \to \llbracket\Gamma_1 \rrbracket \otimes \cdots \otimes \llbracket\Gamma_n \rrbracket$ the morphism that splits $\llbracket\Gamma_1, \ldots, \Gamma_n \rrbracket$ into $\llbracket\Gamma_1 \rrbracket \otimes \cdots \otimes \llbracket\Gamma_n \rrbracket$, and by $\operatorname{jn}_{\Gamma_1;\ldots;\Gamma_n}$ the corresponding inverse. Given a context $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta$ we denote by $\operatorname{exch}_{\Gamma,\underline{x}:\mathbb{A},y:\mathbb{B},\Delta} : \llbracket\Gamma,x:\mathbb{A},y:\mathbb{B},\Delta \rrbracket \to \llbracket\Gamma,y:\mathbb{B},x:\mathbb{A},\Delta \rrbracket$ the morphism corresponding to the permutation of the variable $x:\mathbb{A}$ with $y:\mathbb{B}$. Whenever convenient we will drop variable names in the subscripts of $\operatorname{sp}, \operatorname{jn}, \operatorname{and} \operatorname{exch}$. For a context $E \in \operatorname{Sf}(\Gamma_1,\ldots,\Gamma_n)$ the morphism $\operatorname{sh}_E : \llbracket \rrbracket \to \llbracket\Gamma_1, \ldots, \Gamma_n \rrbracket$ denotes the corresponding shuffling morphism.

For every operation symbol $f : \mathbb{A}_1, \ldots, \mathbb{A}_n \to \mathbb{A}$ in Σ we postulate an interpretation $\llbracket f \rrbracket : \llbracket \mathbb{A}_1 \rrbracket \otimes \cdots \otimes \llbracket \mathbb{A}_n \rrbracket \to \llbracket \mathbb{A} \rrbracket$ as a C-morphism. The interpretation of judgements is defined by induction over the structure of judgement derivation according to the rules in Fig. 2.

$$\begin{array}{ll} \displaystyle \frac{\Gamma_{i} \rhd v_{i} : \mathbb{A}_{i} \quad f : \mathbb{A}_{1}, \dots, \mathbb{A}_{n} \to \mathbb{A} \in \Sigma \quad E \in \mathrm{Sf}(\Gamma_{1}; \dots; \Gamma_{n})}{E \rhd f(v_{1}, \dots, v_{n}) : \mathbb{A}} \quad (\mathbf{ax}) \quad \frac{\pi : \mathbb{A} \rhd x : \mathbb{A}}{x : \mathbb{A} \rhd x : \mathbb{A}} \quad (\mathbf{hyp}) \\ \\ \displaystyle \frac{- \rhd * : \mathbb{I}}{- \rhd * : \mathbb{I}} \quad (\mathbb{I}_{i}) \quad & \frac{\Gamma \rhd v : \mathbb{I} \quad \Delta \rhd w : \mathbb{A} \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{E \rhd v \text{ to } * \cdot w : \mathbb{A}} \quad (\mathbb{I}_{e}) \\ \\ \displaystyle \frac{\Gamma \rhd v : \mathbb{A} \quad \Delta \rhd w : \mathbb{B} \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{E \rhd v \otimes w : \mathbb{A} \otimes \mathbb{B}} \quad (\otimes_{i}) \\ \\ \displaystyle \frac{\Gamma \rhd v : \mathbb{A} \otimes \mathbb{B} \quad \Delta, x : \mathbb{A}, y : \mathbb{B} \rhd w : \mathbb{C} \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{E \rhd \operatorname{pm} v \text{ to } x \otimes y \cdot w : \mathbb{C}} \quad (\otimes_{e}) \\ \\ \displaystyle \frac{\Gamma, x : \mathbb{A} \rhd v : \mathbb{B}}{\Gamma \rhd \lambda x : \mathbb{A}, v : \mathbb{A} \multimap \mathbb{B}} \quad (\multimap_{i}) \quad & \frac{\Gamma \rhd v : \mathbb{A} \multimap \mathbb{B} \quad \Delta \rhd w : \mathbb{A} \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{E \rhd v w : \mathbb{B}} \quad (\multimap_{e}) \end{array}$$

Figure 1 Term formation rules for linear λ -calculus.

As detailed in [4, 27, 28], linear λ -calculus comes equipped with a class of equations (Fig. 3), specifically *equations-in-context* $\Gamma \triangleright v = w : \mathbb{A}$, that corresponds to the axiomatics of autonomous categories. As usual, we omit the context and typing information of the

$$\begin{split} \frac{\llbracket\Gamma_i \rhd v_i : \mathbb{A}_i \rrbracket = m_i \quad f : \mathbb{A}_1, \dots, \mathbb{A}_n \to \mathbb{A} \in \Sigma \quad E \in \mathrm{Sf}(\Gamma_1 \dots \Gamma_n)}{\llbracketE \rhd f(v_1, \dots, v_n) : \mathbb{A} \rrbracket = \llbracketf \rrbracket \cdot (m_1 \otimes \dots \otimes m_n) \cdot \mathrm{sp}_{\Gamma_1; \dots; \Gamma_n} \cdot \mathrm{sh}_E} & \boxed{\llbracketx : \mathbb{A} \rhd x : \mathbb{A} \rrbracket = \mathrm{id}_{\llbracket\mathbb{A}} \rrbracket} \\ \\ \frac{\llbracket\Gamma \rhd v : \mathbb{I} \rrbracket = \mathrm{id}_{\llbracket\mathbb{I}}}{\llbracket - \rhd * : \mathbb{I} \rrbracket = \mathrm{id}_{\llbracket\mathbb{I}}} & \underbrace{\llbracket\Gamma \rhd v : \mathbb{I} \rrbracket = m \quad \llbracket\Delta \rhd w : \mathbb{A} \rrbracket = n \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{\llbracketE \rhd v \text{ to } * . w : \mathbb{A} \rrbracket = n \cdot \lambda \cdot (m \otimes \mathrm{id}) \cdot \mathrm{sp}_{\Gamma; \Delta} \cdot \mathrm{sh}_E} \\ \\ \frac{\llbracket\Gamma \rhd v : \mathbb{A} \rrbracket = m \quad \llbracket\Delta \rhd w : \mathbb{B} \rrbracket = n \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{\llbracketE \rhd v \otimes w : \mathbb{A} \otimes \mathbb{B} \rrbracket = (m \otimes n) \cdot \mathrm{sp}_{\Gamma; \Delta} \cdot \mathrm{sh}_E} & \underbrace{\llbracket\Gamma, x : \mathbb{A} \rhd v : \mathbb{B} \rrbracket = m}{\llbracket\Gamma \rhd v : \mathbb{A} \otimes \mathbb{B} \rrbracket = (m \otimes n) \cdot \mathrm{sp}_{\Gamma; \Delta} \cdot \mathrm{sh}_E} & \underbrace{\llbracket\Gamma \rhd \lambda x : \mathbb{A} \cdot v : \mathbb{B} \rrbracket = m}{\llbracket\Gamma \rhd v : \mathbb{A} \otimes \mathbb{B} \rrbracket = m \quad \llbracket\Delta, x : \mathbb{A}, y : \mathbb{B} \rhd w : \mathbb{C} \rrbracket = n \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{\llbracketE \rhd \operatorname{pm} v \text{ to } x \otimes y \cdot w : \mathbb{C} \rrbracket = n \cdot \mathrm{jn}_{\Delta;\mathbb{A};\mathbb{B}} \cdot \alpha \cdot \mathrm{sw} \cdot (m \otimes \mathrm{id}) \cdot \mathrm{sp}_{\Gamma;\Delta} \cdot \mathrm{sh}_E} \\ \\ \\ \frac{\llbracket\Gamma \rhd v : \mathbb{A} \longrightarrow \mathbb{B} \rrbracket = m \quad \llbracket\Delta \rhd w : \mathbb{A} \rrbracket = m \quad \llbracket\Delta \rhd w : \mathbb{A} \rrbracket = n \quad E \in \mathrm{Sf}(\Gamma; \Delta)}{\llbracketE \rhd \operatorname{pm} v \text{ to } x \otimes y \cdot w : \mathbb{C} \rrbracket = n \cdot \mathrm{jn}_{\Delta;\mathbb{A};\mathbb{B}} \cdot \alpha \cdot \mathrm{sw} \cdot (m \otimes \mathrm{id}) \cdot \mathrm{sp}_{\Gamma;\Delta} \cdot \mathrm{sh}_E} \\ \end{array}$$

Figure 2 Judgement interpretation on an autonomous category C.

equations in Fig. 3, which can be reconstructed in the usual way.

pm $v \otimes w$ to $x \otimes y.$ u	=	u[v/x,w/y]	
pm v to $x\otimes y.$ $u[x\otimes y/z]$	=	u[v/z]	$(\lambda x:\mathbb{A}.\ v)\ w = v[w/x]$
* to *.v	=	v	$\lambda x : \mathbb{A}.(v x) = v$
$v \; { m to} \; * \; . \; w[*/z]$	=	w[v/z]	(b) Higher-order structure
· · · · · ·			

(a) Monoidal structure

 $\begin{array}{rcl} u[v \text{ to } * \,.\, w/z] &=& v \text{ to } * \,.\, u[w/z] \\ u[\texttt{pm} \ v \text{ to } x \otimes y. \ w/z] &=& \texttt{pm} \ v \text{ to } x \otimes y. \ u[w/z] \end{array}$

(c) Commuting conversions

Figure 3 Equations corresponding to the axiomatics of autonomous categories.

▶ **Theorem 3.** The equations presented in Fig. 3 are sound w.r.t. judgement interpretation. Specifically if $\Gamma \triangleright v = w : \mathbb{A}$ is one of the equations in Fig. 3 then $\llbracket \Gamma \triangleright v : \mathbb{A} \rrbracket = \llbracket \Gamma \triangleright w : \mathbb{A} \rrbracket$.

▶ **Definition 4** (Linear λ -theories). Consider a tuple (G, Σ) consisting of a class G of ground types and a class Σ of sorted operation symbols. A linear λ -theory $((G, \Sigma), Ax)$ is a triple such that Ax is a class of equations-in-context over linear λ -terms built from (G, Σ) .

The elements of Ax are called axioms (of the theory). Let Th(Ax) be the smallest congruence that contains Ax, the equations listed in Fig. 3, and that is closed under the exchange and substitution rules. We call the elements of Th(Ax) theorems (of the theory).

▶ Definition 5 (Models of linear λ -theories). Consider a linear λ -theory $((G, \Sigma), Ax)$ and an autonomous category C. Suppose that for each $X \in G$ we have an interpretation [X] that is a C-object and analogously for the operation symbols. This interpretation structure is a model of the theory if all axioms are satisfied by the interpretation.

Next let us turn our attention to the correspondence between linear λ -calculus and autonomous categories, established via soundness, completeness, and internal language theorems. Despite the proofs of such theorems already being detailed in [27, 4, 28], we

decided to briefly sketch them below to render the presentation of some of our own results self-contained.

▶ **Theorem 6** (Soundness & Completeness). Consider a linear λ -theory T. An equation $\Gamma \triangleright v = w : A$ is a theorem of T iff it is satisfied by all models of the theory.

Proof sketch. Soundness follows by induction over the rules that define Th(Ax) (Definition 4) and by Theorem 3. Completeness is based on the idea of a Lindenbaum-Tarski algebra: it follows from building the *syntactic category* Syn(T) of T (also known as term model), showing that it possesses an autonomous structure and also that equality $[\Gamma \triangleright v : \mathbb{A}] = [\Gamma \triangleright w : \mathbb{A}]$ in the syntactic category is equivalent to provability $\Gamma \triangleright v = w : \mathbb{A}$ in the theory.

The syntactic category of T has as objects the types of T and as morphisms $\mathbb{A} \to \mathbb{B}$ the equivalence classes (w.r.t. provability) of terms v for which we can derive $x : \mathbb{A} \rhd v : \mathbb{B}$.

Next let us focus on the topic of internal languages, for which the following result is quite useful.

▶ **Theorem 7.** Consider a linear λ -theory T and a model of T on an autonomous category C. The model induces a functor $F : Syn(T) \rightarrow C$ that (strictly) preserves the autonomous structure.

Proof sketch. Consider a model of T on a category C. Then for any judgement $x : \mathbb{A} \triangleright v : \mathbb{B}$, the induced functor F sends the equivalence class [v] into $[x : \mathbb{A} \triangleright v : \mathbb{B}]$.

An autonomous category C induces a linear λ -theory Lang(C) whose ground types $X \in G$ are the objects of C and whose signature Σ of operation symbols consists of all the morphisms in C plus certain isomorphisms that we describe in (2). The axioms of Lang(C) are all the equations satisfied by the obvious interpretation in C. In order to explicitly distinguish the autonomous structure of C from the type structure of Lang(C) let us denote the tensor of C by $\hat{\otimes}$, the unit by $\hat{\mathbb{I}}$, and the exponential by $\widehat{-\infty}$. Consider then the following map on types:

$$i(\mathbb{I}) = \widehat{\mathbb{I}} \qquad i(X) = X \qquad i(\mathbb{A} \otimes \mathbb{B}) = i(\mathbb{A}) \,\widehat{\otimes} \, i(\mathbb{B}) \qquad i(\mathbb{A} \multimap \mathbb{B}) = i(\mathbb{A}) \,\widehat{\multimap} \, i(\mathbb{B}) \tag{2}$$

For each type A we add an isomorphism $A \simeq i(A)$ to the theory Lang(C).

▶ **Theorem 8** (Internal language). For every autonomous category C there exists an equivalence of categories $Syn(Lang(C)) \simeq C$.

Proof sketch. By construction, we have an interpretation of Lang(C) in C which behaves as the identity for operation symbols and ground types. This interpretation is a model of Lang(C) on C and by Theorem 7 we obtain a functor $\text{Syn}(\text{Lang}(C)) \rightarrow C$. The functor in the opposite direction behaves as the identity on objects and sends a C-morphism f into [f(x)]. The equivalence of categories is then shown by using the aforementioned isomorphisms which connect the type constructors of Lang(C) with the autonomous structure of C.

3 From equations to *V*-equations

We now extend the results of the previous section to the setting of \mathcal{V} -equations.

3.1 A \mathcal{V} -equational deductive system

Let \mathcal{V} denote a commutative and unital quantale, $\otimes : \mathcal{V} \times \mathcal{V} \to \mathcal{V}$ the corresponding binary operation, and k the corresponding unit [32]. As mentioned in the introduction, \mathcal{V} induces the notion of a \mathcal{V} -equation, *i.e.* an equation $t =_q s$ labelled by an element q of \mathcal{V} . This subsection explores this concept by introducing a \mathcal{V} -equational deductive system for linear λ -calculus and a notion of a linear $\mathcal{V}\lambda$ -theory.

Let us start by recalling two definitions concerning ordered structures [14, 16] and then explain their relevance to our work.

▶ **Definition 9.** Consider a complete lattice L. For every $x, y \in L$ we say that y is way-below x (in symbols, $y \ll x$) if for every subset $X \subseteq L$ whenever $x \leq \bigvee X$ there exists a finite subset $A \subseteq X$ such that $y \leq \bigvee A$. The lattice L is called continuous iff for every $x \in L$,

$$x = \bigvee \{ y \mid y \in L \text{ and } y \ll x \}$$

▶ **Definition 10.** Let *L* be a complete lattice. A basis *B* of *L* is a subset $B \subseteq L$ such that for every $x \in L$ the set $B \cap \{y \mid y \in L \text{ and } y \ll x\}$ is directed and has *x* as the least upper bound.

From now on we assume that the underlying lattice of \mathcal{V} is continuous and has a basis B which is closed under finite joins, the multiplication of the quantale \otimes and contains the unit k. These assumptions will allow us to work *only* with a specified subset of \mathcal{V} -equations chosen *e.g.* for computational reasons, such as the *finite* representation of values $q \in \mathcal{V}$.

▶ **Example 11.** The Boolean quantale $((\{0 \le 1\}, \lor), \otimes := \land)$ is *finite* and thus continuous [14]. Since it is continuous, $\{0, 1\}$ itself is a basis for the quantale that satisfies the conditions above. For the Gödel t-norm [11] $(([0, 1], \lor), \otimes := \land)$, the way-below relation is the strictly-less relation < with the exception that 0 < 0. A basis for the underlying lattice that satisfies the conditions above is the set $\mathbb{Q} \cap [0, 1]$. Note that, unlike real numbers, rationals numbers always have a finite representation. For the metric quantale (also known as Lawvere quantale) $(([0, \infty], \land), \otimes := +)$, the way-below relation corresponds to the *strictly greater* relation with $\infty > \infty$, and a basis for the underlying lattice that satisfies the conditions above is the set of extended non-negative rational numbers. The latter also serves as basis for the ultrametric quantale ($([0, \infty], \land), \otimes := \max$).

We also assume that \mathcal{V} is *integral*, *i.e.* that the unit k is the top element of \mathcal{V} . This will allow us to establish a smoother theory of \mathcal{V} -equations, whilst still covering *e.g.* all the examples above. This assumption is common in quantale theory [39].

Recall the term formation rules of linear λ -calculus from Fig. 1. A \mathcal{V} -equation-in-context is an expression $\Gamma \rhd v =_q w : \mathbb{A}$ with $q \in B$ (the basis of \mathcal{V}), $\Gamma \rhd v : \mathbb{A}$ and $\Gamma \rhd w : \mathbb{A}$. Let \top be the top element in \mathcal{V} . An equation-in-context $\Gamma \rhd v = w : \mathbb{A}$ now denotes the particular case in which both $\Gamma \rhd v =_{\top} w : \mathbb{A}$ and $\Gamma \rhd w =_{\top} v : \mathbb{A}$. For the case of the Boolean quantale, \mathcal{V} -equations are labelled by $\{0,1\}$. We will see that $\Gamma \rhd v =_1 w : \mathbb{A}$ can be treated as an inequation $\Gamma \rhd v \leq w : \mathbb{A}$, whilst $\Gamma \rhd v =_0 w : \mathbb{A}$ corresponds to a trivial \mathcal{V} -equation, *i.e.* a \mathcal{V} -equation that always holds. For the Gödel t-norm, we can choose $\mathbb{Q} \cap [0,1]$ as basis and then obtain what we call *fuzzy inequations*. For the metric quantale, we can choose the set of extended non-negative rational numbers as basis and then obtain *metric equations* in the spirit of [29, 30]. Similarly, by choosing the ultrametric quantale (($[0, \infty], \wedge$), $\otimes := \max$) with the set of extended non-negative rational numbers as basis we obtain what we call *ultrametric equations*.

▶ Definition 12 (Linear $\mathcal{V}\lambda$ -theories). Consider a tuple (G, Σ) consisting of a class G of ground types and a class of sorted operation symbols $f : \mathbb{A}_1, \ldots, \mathbb{A}_n \to \mathbb{A}$ with $n \ge 1$. A linear $\mathcal{V}\lambda$ -theory $((G, \Sigma), Ax)$ is a tuple such that Ax is a class of \mathcal{V} -equations-in-context over linear λ -terms built from (G, Σ) .

 $\begin{array}{ll} \overline{v=_{\top}v} \ (\text{refl}) & \frac{v=_{q}w}{v=_{q\otimes r}u} \frac{w=_{r}u}{w} \ (\text{trans}) & \frac{v=_{q}w}{v=_{r}w} \frac{r\leq q}{v=_{r}w} \ (\text{weak}) \\ & \frac{\forall r\ll q, v=_{r}w}{v=_{q}w} \ (\text{arch}) & \frac{\forall i\leq n. \ v=_{q_{i}}w}{v=_{\sqrt{q_{i}}w}} \ (\text{join}) \\ & \frac{\forall i\leq n. \ v_{i}=_{q_{i}}w_{i}}{f(v_{1},\ldots,v_{n})=_{\otimes q_{i}}f(w_{1},\ldots,w_{n})} & \frac{v=_{q}w}{v\otimes v'=_{r}w'} \\ & \frac{v=_{q}w}{v\otimes v'=_{q\otimes r}w\otimes w'} \\ & \frac{v=_{q}w}{v \ to \ x. \ v'=_{q\otimes r}w} & \frac{v=_{q}w}{v'=_{r}w'} \\ & \frac{v=_{q}w}{v \ to \ x. \ v'=_{q\otimes r}w \ to \ x. \ w'}} & \frac{v=_{q}w}{\lambda x: \mathbb{A}. \ v=_{q}\lambda x: \mathbb{A}. \ w} & \frac{v=_{q}w}{v'=_{r}w'} \\ & \frac{v=_{q}w}{v(v'=_{r}w')} & \frac{v=_{q}w}{v(v'=_{r}w')} \\ & \frac{v=_{q}w}{v(v'=_{q\otimes r}w)} \otimes \frac{v=_{q}w}{v'=_{q\otimes r}w'} \\ & \frac{v=_{q}w}{\Delta v=_{q}w: \mathbb{A}} & \frac{\Delta \in perm(\Gamma)}{v(v'/x]=_{q\otimes r}w[w'/x]} \end{array}$

Figure 4 *V*-congruence rules.

The elements of Ax are the axioms of the theory. Let Th(Ax) be the smallest class that contains Ax and that is closed under the rules of Fig. 3 and of Fig. 4 (as usual we omit the context and typing information). The elements of Th(Ax) are the theorems of the theory.

Let us examine the rules in Fig. 4 in more detail. They can be seen as a generalisation of the notion of a congruence. The rules (refl) and (trans) are a generalisation of equality's reflexivity and transitivity. Rule (weak) encodes the principle that the higher the label in the \mathcal{V} -equation, the 'tighter' is the relation between the two terms in the \mathcal{V} -equation. In other words, $v =_r w$ is subsumed by $v =_q w$, for $r \leq q$. This can be seen clearly *e.g.* with the metric quantale by reading $v =_q w$ as "the terms v and w are *at most* at distance q from each other" (recall that in the metric quantale the usual order is reversed, *i.e.* $\leq := \geq_{[0,\infty]}$). (arch) is essentially a generalisation of the Archimedean rule in [29, 30]. It says that if $v =_r w$ for all approximations r of q then it is also the case that $v =_q w$. (join) says that deductions are closed under finite joins, and in particular it is always the case that $v =_{\perp} w$. All other rules correspond to a generalisation of *compatibility* to a \mathcal{V} -equational setting.

The reader may have noticed that the rules in Fig. 4 do not contain a \mathcal{V} -generalisation of symmetry w.r.t. standard equality. Such a generalisation would be:

$$\frac{v =_q u}{w =_q u}$$

This rule is not present in Fig. 4 because in some quantales \mathcal{V} it forces too many \mathcal{V} -equations. For example, in the Boolean quantale the condition $v \leq w$ would automatically entail $w \leq v$ (due to symmetry); in fact, for this particular case symmetry forces the notion of inequation to collapse into the classical notion of equation. On the other hand, symmetry is desirable in the (ultra)metric case because (ultra)metrics need to respect the symmetry equation [16].

▶ **Definition 13** (Symmetric linear $V\lambda$ -theories). A symmetric linear $V\lambda$ -theory is a linear $V\lambda$ -theory whose set of theorems is closed under symmetry.

In Appendix A we further explore how specific families of quantales are reflected in the \mathcal{V} -equational system here introduced, and briefly compare the latter to metric algebra [29, 30].

3.2 Semantics of *V***-equations**

In this subsection we set the necessary background for presenting a sound and complete class of models for (symmetric) linear $\mathcal{V}\lambda$ -theories. We start by recalling basics concepts of \mathcal{V} -categories, which are central in a field initiated by Lawvere in [24] and can be intuitively seen as generalised metric spaces [38, 17, 39]. As we will see, \mathcal{V} -categories provide structure to suitably interpret \mathcal{V} -equations.

▶ Definition 14. A (small) \mathcal{V} -category is a pair (X, a) where X is a class (set) and a : $X \times X \rightarrow \mathcal{V}$ is a function that satisfies:

 $k \le a(x, x)$ and $a(x, y) \otimes a(y, z) \le a(x, z)$ $(x, y, z \in X)$

For two \mathcal{V} -categories (X, a) and (Y, b), a \mathcal{V} -functor $f : (X, a) \to (Y, b)$ is a function $f : X \to Y$ that satisfies the inequality $a(x, y) \leq b(f(x), f(y))$ for all $x, y \in X$.

Small \mathcal{V} -categories and \mathcal{V} -functors form a category which we denote by \mathcal{V} -Cat. A \mathcal{V} -category (X, a) is called *symmetric* if a(x, y) = a(y, x) for all $x, y \in X$. We denote by \mathcal{V} -Cat_{sym} the full subcategory of \mathcal{V} -Cat whose objects are symmetric. Every \mathcal{V} -category carries a natural order defined by $x \leq y$ whenever $k \leq a(x, y)$. A \mathcal{V} -category is called *separated* if its natural order is anti-symmetric. We denote by \mathcal{V} -Cat_{sep} the full subcategory of \mathcal{V} -Cat whose objects are separated.

▶ **Example 15.** For \mathcal{V} the Boolean quantale, \mathcal{V} -Cat_{sep} is the category Pos of partially ordered sets and monotone maps; \mathcal{V} -Cat_{sym,sep} is simply the category Set of sets and functions. For \mathcal{V} the metric quantale, \mathcal{V} -Cat_{sym,sep} is the category Met of extended metric spaces and non-expansive maps. In what follows we omit the qualifier 'extended' in 'extended (ultra)metric spaces'. For \mathcal{V} the ultrametric quantale, \mathcal{V} -Cat_{sym,sep} is the category of ultrametric spaces and non-expansive maps.

The inclusion functor $\mathcal{V}\text{-}\mathsf{Cat}_{\mathsf{sep}} \hookrightarrow \mathcal{V}\text{-}\mathsf{Cat}$ has a left adjoint [17]. It is constructed first by defining the equivalence relation $x \sim y$ whenever $x \leq y$ and $y \leq x$ (for \leq the natural order introduced earlier). Then this relation induces the separated $\mathcal{V}\text{-}category$ $(X/_{\sim}, \tilde{a})$ where \tilde{a} is defined as $\tilde{a}([x], [y]) = a(x, y)$ for every $[x], [y] \in X/_{\sim}$. The left adjoint of the inclusion functor $\mathcal{V}\text{-}\mathsf{Cat}_{\mathsf{sep}} \hookrightarrow \mathcal{V}\text{-}\mathsf{Cat}$ sends every $\mathcal{V}\text{-}category$ (X, a) to $(X/_{\sim}, \tilde{a})$. This quotienting construct preserves symmetry, and therefore we automatically obtain the following result.

▶ Theorem 16. The inclusion functor \mathcal{V} -Cat_{sym,sep} $\hookrightarrow \mathcal{V}$ -Cat_{sym} has a left adjoint.

Next, we recall notions of enriched category theory [19] instantiated into the setting of autonomous categories enriched over \mathcal{V} -categories. We will use the enriched structure to give semantics to \mathcal{V} -equations between linear λ -terms. First, note that every category \mathcal{V} -Cat is autonomous with the tensor $(X, a) \otimes (Y, b) := (X \times Y, a \otimes b)$ where $a \otimes b$ is defined as $(a \otimes b)((x, y), (x', y')) = a(x, x') \otimes b(y, y')$ and the set of \mathcal{V} -functors \mathcal{V} -Cat((X, a), (Y, b)) equipped with the map $(f, g) \mapsto \bigwedge_{x \in X} b(f(x), g(x))$.

▶ **Theorem 17.** The categories \mathcal{V} -Cat_{sym}, \mathcal{V} -Cat_{sym,sep} inherit the autonomous structure of \mathcal{V} -Cat whenever \mathcal{V} is integral.

Since we assume that \mathcal{V} is integral, this last theorem allows us to formally define the notion of categories enriched over \mathcal{V} -categories using [19].

▶ **Definition 18.** A category C is \mathcal{V} -Cat-enriched (or simply, a \mathcal{V} -Cat-category) if for all C-objects X and Y the hom-set C(X, Y) is a \mathcal{V} -category and if the composition of C-morphisms,

 (\cdot) : $C(X,Y) \otimes C(Y,Z) \longrightarrow C(X,Z)$

is a \mathcal{V} -functor. Given two \mathcal{V} -Cat-categories C and D and a functor $F : \mathsf{C} \to \mathsf{D}$, we call F a \mathcal{V} -Cat-functor if for all C -objects X and Y the map $F_{X,Y} : \mathsf{C}(X,Y) \to \mathsf{D}(FX,F,Y)$ is a \mathcal{V} -functor. An adjunction $\mathsf{C} : F \dashv G : \mathsf{D}$ is called \mathcal{V} -Cat-enriched if the underlying functors F and G are \mathcal{V} -Cat-functors and if for all objects $X \in |\mathsf{C}|$ and $Y \in |\mathsf{D}|$ there exists a \mathcal{V} -isomorphism $\mathsf{D}(FX,Y) \simeq \mathsf{C}(X,GY)$ natural in X and Y.

If C is a \mathcal{V} -Cat-category then $C \times C$ is also a \mathcal{V} -Cat-category via the tensor operation \otimes in \mathcal{V} -Cat. We take advantage of this fact in the following definition.

▶ Definition 19. A \mathcal{V} -Cat-enriched autonomous category C is an autonomous and \mathcal{V} -Catcategory C such that the bifunctor \otimes : C × C → C is a \mathcal{V} -Cat-functor and the adjunction $(-\otimes X) \dashv (X \multimap -)$ is a \mathcal{V} -Cat-adjunction.

▶ **Example 20.** Recall that $Pos \simeq \mathcal{V}\text{-}Cat_{sep}$ when \mathcal{V} is the Boolean quantale. According to Theorem 17 the category Pos is autonomous. It follows by general results that the category is Pos-enriched [6]. It is also easy to see that its tensor is Pos-enriched and that the adjunction $(- \otimes X) \dashv (X \multimap -)$ is Pos-enriched. Therefore, Pos is an instance of Definition 19. Note also that Set $\simeq \mathcal{V}\text{-}Cat_{sym,sep}$ for \mathcal{V} the Boolean quantale and that Set is an instance of Definition 19.

Recall that $Met \simeq \mathcal{V}\text{-}Cat_{sym,sep}$ when \mathcal{V} is the metric quantale. Thus, the category Met is autonomous (Theorem 17) and Met-enriched [6]. It follows as well from routine calculations that its tensor is Met-enriched and that the adjunction $(- \otimes X) \dashv (X \multimap -)$ is Met-enriched. Therefore Met is an instance of Definition 19. An analogous reasoning tells that the category of ultrametric spaces (enriched over itself) is also an instance of Definition 19.

Finally, recall the interpretation of linear λ -terms on an autonomous category C (Section 2) and assume that C is \mathcal{V} -Cat-enriched. Then we say that a \mathcal{V} -equation $\Gamma \triangleright v =_q w : \mathbb{A}$ is satisfied by this interpretation if $a(\llbracket\Gamma \triangleright v : \mathbb{A}\rrbracket, \llbracket\Gamma \triangleright w : \mathbb{A}\rrbracket) \ge q$ where $a : \mathsf{C}(\llbracket\Gamma\rrbracket, \llbracket\mathbb{A}\rrbracket) \times \mathsf{C}(\llbracket\Gamma\rrbracket, \llbracket\mathbb{A}\rrbracket) \to \mathcal{V}$ is the underlying function of the \mathcal{V} -category $\mathsf{C}(\llbracket\Gamma\rrbracket, \llbracket\mathbb{A}\rrbracket)$.

▶ **Theorem 21.** The rules listed in Fig. 3 and Fig. 4 are sound for \mathcal{V} -Cat-enriched autonomous categories C. Specifically, if $\Gamma \rhd v =_q w$: A results from the rules in Fig. 3 and Fig. 4 then $a(\llbracket \Gamma \rhd v : A\rrbracket, \llbracket \Gamma \rhd w : A\rrbracket) \ge q$.

Proof of Theorem 21. Let us focus first on the equations listed in Fig. 3. Recall that an equation $\Gamma \triangleright v = w$: A abbreviates the \mathcal{V} -equations $\Gamma \triangleright v =_{\top} w$: A and $\Gamma \triangleright w =_{\top} v$: A. Moreover, we already know that the equations listed in Fig. 3 are sound for autonomous categories, specifically if v = w is an equation of Fig. 3 then $\llbracket v \rrbracket = \llbracket w \rrbracket$ in C (Theorem 3). Thus, by the definition of a \mathcal{V} -category and by the assumption of \mathcal{V} being integral $(k = \top)$ we obtain $a(\llbracket v \rrbracket, \llbracket w \rrbracket) \ge k = \top$ and $a(\llbracket w \rrbracket, \llbracket v \rrbracket) \ge k = \top$.

Let us now focus on the rules listed in Fig. 4. The first three rules follow from the definition of a \mathcal{V} -category and the transitivity property of \leq . Rule (arch) follows from the continuity of \mathcal{V} , specifically from the fact that q is the *least* upper bound of all elements r that are way-below q. Rule (join) follows from the definition of least upper bound. The remaining rules follow from the definition of the tensor functor \otimes in \mathcal{V} -Cat, the fact that C is \mathcal{V} -Cat-enriched, $\otimes : C \times C \to C$ is a \mathcal{V} -Cat-functor, and the fact that $(- \otimes X) \dashv (X \multimap -)$ is

a \mathcal{V} -Cat-adjunction. For example, for the sixth rule we reason as follows:

$$\begin{aligned} a(\llbracket f(v_1, \dots, v_n) \rrbracket, \llbracket f(w_1, \dots, w_n) \rrbracket) \\ &= a(\llbracket f \rrbracket \cdot (\llbracket v_1 \rrbracket \otimes \dots \otimes \llbracket v_n \rrbracket) \cdot \mathsf{sp}_{\Gamma_1; \dots; \Gamma_n} \cdot \mathsf{sh}_E, \llbracket f \rrbracket \cdot (\llbracket w_1 \rrbracket \otimes \dots \otimes \llbracket w_n \rrbracket) \cdot \mathsf{sp}_{\Gamma_1; \dots; \Gamma_n} \cdot \mathsf{sh}_E) \\ &\geq a(\llbracket f \rrbracket \cdot (\llbracket v_1 \rrbracket \otimes \dots \otimes \llbracket v_n \rrbracket), \llbracket f \rrbracket \cdot (\llbracket w_1 \rrbracket \otimes \dots \otimes \llbracket w_n \rrbracket)) \\ &\geq a(\llbracket v_1 \rrbracket \otimes \dots \otimes \llbracket v_n \rrbracket), (\llbracket w_1 \rrbracket \otimes \dots \otimes \llbracket w_n \rrbracket) \\ &\geq a(\llbracket v_1 \rrbracket, \llbracket w_1 \rrbracket) \otimes \dots \otimes a(\llbracket v_n \rrbracket, \llbracket w_n \rrbracket) \\ &\geq a(\llbracket v_1 \rrbracket, \llbracket w_1 \rrbracket) \otimes \dots \otimes a(\llbracket v_n \rrbracket, \llbracket w_n \rrbracket) \end{aligned}$$

where the second step follows from the fact that $sp_{\Gamma_1;...;\Gamma_n} \cdot sh_E$ is a morphism in C and that C is \mathcal{V} -Cat-enriched. The third step follows from an analogous reasoning. The fourth step follows from the fact that $\otimes : C \times C \to C$ is a \mathcal{V} -Cat-functor. The last step follows from the premise of the rule in question. As another example, the proof for the substitution rule proceeds similarly:

$$\begin{split} &a(\llbracket v[v'/x]\rrbracket, \llbracket w[w'/x]\rrbracket) \\ &= a(\llbracket v\rrbracket \cdot \mathsf{jn}_{\Gamma,\mathbb{A}} \cdot (\mathsf{id} \otimes \llbracket v'\rrbracket) \cdot \mathsf{sp}_{\Gamma;\Delta}, \llbracket w\rrbracket \cdot \mathsf{jn}_{\Gamma,\mathbb{A}} \cdot (\mathsf{id} \otimes \llbracket w'\rrbracket) \cdot \mathsf{sp}_{\Gamma;\Delta}) \\ &\geq a(\llbracket v\rrbracket \cdot \mathsf{jn}_{\Gamma,\mathbb{A}} \cdot (\mathsf{id} \otimes \llbracket v'\rrbracket), \llbracket w\rrbracket \cdot \mathsf{jn}_{\Gamma,\mathbb{A}} \cdot (\mathsf{id} \otimes \llbracket w'\rrbracket)) \\ &\geq a(\mathsf{id} \otimes \llbracket v'\rrbracket, \mathsf{id} \otimes \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket \cdot \mathsf{jn}_{\Gamma,\mathbb{A}}, \llbracket w\rrbracket \cdot \mathsf{jn}_{\Gamma,\mathbb{A}}) \\ &\geq a(\mathsf{id} \otimes \llbracket v'\rrbracket, \mathsf{id} \otimes \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket, \llbracket w\rrbracket) \\ &\geq a(\mathsf{id} \otimes \llbracket v'\rrbracket, \mathsf{id} \otimes \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket, \llbracket w\rrbracket) \\ &\geq a(\mathsf{id} \otimes \llbracket v'\rrbracket, \mathsf{id} \otimes \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket, \llbracket w\rrbracket) \\ &\geq a(\mathsf{id}, \mathsf{id}) \otimes a(\llbracket v'\rrbracket, \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket, \llbracket w\rrbracket) \\ &\geq a(\mathsf{id}, \mathsf{id}) \otimes a(\llbracket v'\rrbracket, \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket, \llbracket w\rrbracket) \\ &= a(\llbracket v'\rrbracket, \llbracket w'\rrbracket) \otimes a(\llbracket v\rrbracket, \llbracket w\rrbracket) \\ &\geq q \otimes r \end{split}$$

The proof for the rule concerning $(-\circ_i)$ additionally requires the following two facts: if a \mathcal{V} -functor $f:(X,a) \to (Y,b)$ is an isomorphism then a(x,x') = b(f(x), f(x')) for all $x, x' \in X$. For a context Γ , the morphism $\mathsf{jn}_{\Gamma;x:\mathbb{A}}: \llbracket\Gamma \rrbracket \otimes \llbracket\mathbb{A} \rrbracket \to \llbracket\Gamma, x:\mathbb{A} \rrbracket$ is an isomorphism in C . The proof for the rule concerning the permutation of variables (exchange) also makes use of the fact that $\llbracket\Delta \rrbracket \to \llbracket\Gamma \rrbracket$ is an isomorphism.

3.3 Soundness, completeness, and internal language

In this subsection we establish a formal correspondence between linear $\mathcal{V}\lambda$ -theories and \mathcal{V} -Cat-enriched autonomous categories, via soundness, completeness, and internal language theorems. A key construct in this correspondence is the quotienting of a \mathcal{V} -category into a *separated* \mathcal{V} -category: we will use it to identify linear λ -terms when generating a syntactic category (from a linear $\mathcal{V}\lambda$ -theory) that satisfies the axioms of autonomous categories. This naturally leads to the following notion of a model for linear $\mathcal{V}\lambda$ -theories.

▶ Definition 22 (Models of linear $V\lambda$ -theories). Consider a linear $V\lambda$ -theory $((G, \Sigma), Ax)$ and a V-Cat_{sep}-enriched autonomous category C. Suppose that for each $X \in G$ we have an interpretation $[\![X]\!]$ as a C-object and analogously for the operation symbols. This interpretation structure is a model of the theory if all axioms in Ax are satisfied by the interpretation.

Another thing that we need to take into account is the size of categories. In Section 2 we did not assume that autonomous categories should be locally small. In particular linear λ -theories are able to generate non-(locally small) categories. Now we need to be stricter because \mathcal{V} -Cat_{sep}-enriched autonomous categories are always locally small (recall the

definition of $\mathcal{V}\text{-}\mathsf{Cat}_{\mathsf{sep}}$). Thus for two types \mathbb{A} and \mathbb{B} of a $\mathcal{V}\lambda$ -theory T, consider the class $\mathsf{Values}(\mathbb{A},\mathbb{B})$ of values v such that $x : \mathbb{A} \rhd v : \mathbb{B}$. We equip $\mathsf{Values}(\mathbb{A},\mathbb{B})$ with the function $a : \mathsf{Values}(\mathbb{A},\mathbb{B}) \times \mathsf{Values}(\mathbb{A},\mathbb{B}) \to \mathcal{V}$ defined by,

$$a(v,w) = \bigvee \{q \mid v =_q w \text{ is a theorem of } T\}$$

It is easy to see that $(Values(\mathbb{A}, \mathbb{B}), a)$ is a (possibly large) \mathcal{V} -category. We then quotient this \mathcal{V} -category into a *separated* \mathcal{V} -category which we suggestively denote by $C(\mathbb{A}, \mathbb{B})$ (as detailed in the proof of the next theorem, $C(\mathbb{A}, \mathbb{B})$ will serve as a hom-object of a syntactic category C generated from a linear $\mathcal{V}\lambda$ -theory). Following the nomenclature of [25], we call T varietal if $C(\mathbb{A}, \mathbb{B})$ is a small \mathcal{V} -category. In the rest of the paper we will only work with varietal theories and locally small categories.

▶ **Theorem 23** (Soundness & Completeness). Consider a varietal $V\lambda$ -theory. A V-equationin-context $\Gamma \triangleright v =_q w : A$ is a theorem iff it holds in all models of the theory.

Proof sketch. Soundness follows by induction over the rules that define the class Th(Ax) (Definition 12) and by Theorem 21. For completeness, we use a strategy similar to the proof of Theorem 6, and take advantage of the quotienting of a \mathcal{V} -category into a separated \mathcal{V} -category. Recall that we assume that the theory is *varietal* and therefore can safely take $C(\mathbb{A}, \mathbb{B})$ to be a small \mathcal{V} -category. Note that the quotienting process identifies all terms $x : \mathbb{A} \rhd v : \mathbb{B}$ and $x : \mathbb{A} \rhd w : \mathbb{B}$ such that $v =_{\top} w$ and $w =_{\top} v$. Such a relation contains the equations-in-context from Fig. 3 and moreover it is straighforward to show that it is compatible with the term formation rules of linear λ -calculus (Fig. 1). So, analogously to Theorem 6 we obtain an autonomous category C whose objects are the types of the language and whose hom-sets are the underlying sets of the \mathcal{V} -categories $C(\mathbb{A}, \mathbb{B})$.

Our next step is to show that the category C has a \mathcal{V} -Cat_{sep}-enriched autonomous structure. We start by showing that the composition map $C(\mathbb{A}, \mathbb{B}) \otimes C(\mathbb{B}, \mathbb{C}) \to C(\mathbb{A}, \mathbb{C})$ is a \mathcal{V} -functor:

$$\begin{aligned} a(([v'], [v]), ([w'], [w])) &= a([v], [w]) \otimes a([v'], [w']) \\ &= a(v, w) \otimes a(v', w') \\ &= \bigvee \{q \mid v =_q w\} \otimes \bigvee \{r \mid v' =_r w'\} \\ &= \bigvee \{q \otimes r \mid v =_q w, v' =_r w'\} \\ &\leq \bigvee \{q \mid v[v'/x] =_q w[w'/x]\} \qquad (A \subseteq B \Rightarrow \bigvee A \leq \bigvee B) \\ &= a(v[v'/x], w[w'/x]) \\ &= a([v[v'/x]], [w[w'/x]]) \\ &= a([v[v'/x]], [w[w'/x]]) \\ &= a([v[v'/x]], [w[w'/x]]) \end{aligned}$$

The fact that $\otimes : \mathsf{C} \times \mathsf{C} \to \mathsf{C}$ is a \mathcal{V} -Cat-functor follows by an analogous reasoning. Next, we need to show that $(- \otimes X) \dashv (X \multimap -)$ is a \mathcal{V} -Cat-adjunction. It is straightforward to show that both functors are \mathcal{V} -Cat-functors, and from a similar reasoning it follows that the isomorphism $\mathsf{C}(\mathbb{B}, \mathbb{A} \multimap \mathbb{C}) \simeq \mathsf{C}(\mathbb{B} \otimes \mathbb{A}, \mathbb{C})$ is a \mathcal{V} -isomorphism.

The final step is to show that if an equation $\Gamma \triangleright v =_q w : \mathbb{A}$ with $q \in B$ is satisfied by C then it is a theorem of the linear $\mathcal{V}\lambda$ -theory. By assumption $a([v], [w]) = a(v, w) = \bigvee \{r \mid v =_r w\} \ge q$. It follows from the definition of the way-below relation that for all $x \in B$ with $x \ll q$ there exists a *finite* set $A \subseteq \{r \mid v =_r w\}$ such that $x \le \bigvee A$. Then by an application of rule (join) in Fig. 4 we obtain $v =_{\bigvee A} w$, and consequently rule (weak) in Fig. 4 provides $v =_x w$ for all $x \ll q$. Finally, by an application of rule (arch) in Fig. 4 we deduce that $v =_q w$ is part of the theory.

Next we establish results that will be key in the proof of the internal language theorem. Let Syn(T) be syntactic category of a linear $\mathcal{V}\lambda$ -theory T, as described in Theorem 23.

▶ **Theorem 24.** Consider a linear $\mathcal{V}\lambda$ -theory T and a model of T on a \mathcal{V} -Cat_{sep}-enriched autonomous category C . The model induces a \mathcal{V} -Cat_{sep}-functor $\operatorname{Syn}(T) \to \mathsf{C}$ that (strictly) preserves the autonomous structure of $\operatorname{Syn}(T)$.

Proof. Consider a model of T over C . Let a denote the underlying function of the hom-(\mathcal{V} -categories) in $\operatorname{Syn}(T)$ and b the underlying function of the hom-(\mathcal{V} -categories) in C . Then note that if [v] = [w] then, by completeness, the equations $v =_{\top} w$ and $w =_{\top} v$ are theorems, which means that $[\![v]\!] = [\![w]\!]$ by the definition of a model and *separability*. This allows us to define a mapping $F : \operatorname{Syn}(T) \to \mathsf{C}$ that sends each type \mathbb{A} to $[\![\mathbb{A}]\!]$ and each morphism [v] to $[\![v]\!]$. The fact that this mapping is an autonomous functor follows from an analogous reasoning to the one used in the proof of Theorem 7. We now need to show that this functor is \mathcal{V} - $\operatorname{Cat}_{\operatorname{sep}}$ -enriched. Recall that $a([v], [w]) = \bigvee\{q \mid v =_q w\}$ and observe that for every $v =_q w$ in the previous quantification we have $b([\![v]\!], [\![w]\!]) \ge q$ (by the definition of a model), which establishes, by the definition of a least upper bound, $a([v], [w]) = \bigvee\{q \mid v =_q w\} \le b([\![v]\!], [\![w]\!])$.

Consider now a \mathcal{V} -Cat_{sep}-enriched autonomous category C. It induces a linear $\mathcal{V}\lambda$ -theory Lang(C) whose ground types and operations symbols are defined as in the case of linear λ -theories (recall Section 2). The axioms of Lang(C) are all the \mathcal{V} -equations-in-context that are satisfied by the obvious interpretation on C.

▶ Theorem 25. The linear $V\lambda$ -theory Lang(C) is varietal.

In conjunction with the proof of Theorem 23, a consequence of this last theorem is that Syn(Lang(C)) is a \mathcal{V} -Cat_{sep}-enriched category. Then we state,

▶ **Theorem 26** (Internal language). For every \mathcal{V} -Cat_{sep}-enriched autonomous category C there exists a \mathcal{V} -Cat_{sep}-equivalence of categories Syn(Lang(C)) \simeq C.

Proof. Let a denote the underlying function of the hom-(\mathcal{V} -categories) in Syn(Lang(C)) and b the underlying function of the hom-(\mathcal{V} -categories) in C. We have, by construction, a model of Lang(C) on C which acts as the identity in the interpretation of ground types and operation symbols. We can then appeal to Theorem 24 to establish a \mathcal{V} -Cat_{sep}-functor Syn(Lang(C)) \rightarrow C. Next, the functor working on the inverse direction behaves as the identity on objects and sends a morphism f into [f(x)]. Let us show that it is \mathcal{V} -Cat_{sep}-enriched. First, observe that if $q \ll b(f,g)$ in C and $q \in B$ then $f(x) =_q g(x)$ is a theorem of Lang(C), due to the fact that \ll entails \leq and by the definition of Lang(C). Using the definition of a basis, we thus obtain $b(f,g) = \bigvee\{q \in B \mid q \ll b(f,g)\} \leq \bigvee\{q \in B \mid f(x) =_q g(x)\} = a([f(x)], [g(x)])$. The equivalence of categories is then shown as in the proof of Theorem 8.

All the results in this section can be extended straightforwardly to the case of *symmetric* linear $\mathcal{V}\lambda$ -theories and \mathcal{V} -Cat_{sym,sep}-enriched autonomous categories.

4 Examples of linear $V\lambda$ -theories and their models

▶ Example 27 (Wait calls). We now return to the example of wait calls and the corresponding metric axioms (1) sketched in the Introduction. Let us build a model over Met for this theory: fix a metric space A, interpret the ground type X as $\mathbb{N} \otimes A$ and the operation symbol wait_n : $X \to X$ as the non-expansive map, $\llbracket \texttt{wait}_n \rrbracket : \mathbb{N} \otimes A \to \mathbb{N} \otimes A$, $(i, a) \mapsto (i + n, a)$.

Since we already know that Met is enriched over itself (recall Definition 19 and Example 20) we only need to show that the axioms in (1) are satisfied by the proposed interpretation. This can be shown via a few routine calculations.

Now, it may be the case that is unnecessary to know the *distance* between the execution time of two programs – instead it suffices to know whether a program finishes its execution *before* another one. This leads us to linear $\mathcal{V}\lambda$ -theories where \mathcal{V} is the Boolean quantale. We call such theories *linear ordered* λ -theories. Recall the language from the Introduction with a single ground type X and the signature of wait calls $\Sigma = {\texttt{wait}_n : X \to X \mid n \in \mathbb{N}}$. Then we adapt the metric axioms (1) to the case of the Boolean quantale by considering instead:

$$\texttt{wait}_{\texttt{O}}(x) = x \qquad \texttt{wait}_{\texttt{n}}(\texttt{wait}_{\texttt{m}}(x)) = \texttt{wait}_{\texttt{n}+\texttt{m}}(x) \qquad \frac{n \leq m}{\texttt{wait}_{\texttt{n}}(x) \leq \texttt{wait}_{\texttt{m}}(x)}$$

where a classical equation v = w is shorthand for $v \le w$ (*i.e.* $v =_1 w$) and $w \le v$ (*i.e.* $w =_1 v$). In the resulting theory we can consider for instance (and omitting types for simplicity) the λ -term that defines the composition of two functions λf . λg . g (f x), which we denote by v, and show that:

$$v \ (\lambda x. \texttt{wait}_1(x)) \leq v \ (\lambda x. \texttt{wait}_1(\texttt{wait}_1(x)))$$

This inequation between higher-order programs arises from the argument $\lambda x. \operatorname{wait}_1(\operatorname{wait}_1(x))$ being costlier than the argument $\lambda x. \operatorname{wait}_1(x)$ – specifically, the former will invoke one more wait call (wait_1) than the latter. Moreover, the inequation entails that for every argument g the execution time of computation v ($\lambda x. \operatorname{wait}_1(x)$) g will always be smaller than that of computation v ($\lambda x. \operatorname{wait}_1(\operatorname{wait}_1(x)$)) g since it invokes one more wait call. Thus in general the inequation tells that costlier programs fed as input to v will result in longer execution times when performing the corresponding computation. In order to build a model for the ordered theory of wait calls, we consider a poset A and define a model over Pos by sending X into $\mathbb{N} \otimes A$ and $\operatorname{wait}_n : X \to X$ to the monotone map $[\operatorname{wait}_n] : \mathbb{N} \otimes A \to \mathbb{N} \otimes A$, $(i, a) \mapsto (i + n, a)$. Since we already know that Pos is enriched over itself (recall Definition 19 and Example 20) we only need to show that the ordered axioms are satisfied by the proposed interpretation. But again, this can be shown via a few routine calculations.

▶ Example 28 (Probabilistic programs). We consider ground types Real, Real⁺, unit and a signature consisting of $\{r : \mathbb{I} \to \text{Real} \mid r \in \mathbb{Q}\} \cup \{r^+ : \mathbb{I} \to \text{Real}^+ \mid r \in \mathbb{Q}_{\geq 0}\} \cup \{r^u : \mathbb{I} \to \text{unit} \mid r \in [0,1] \cap \mathbb{Q}\}$, an operation + of type Real, Real → Real, and sampling functions bernoulli : Real, Real, unit → Real and normal : Real, Real⁺ → Real. Whenever no ambiguities arise, we drop the superscripts in r^u and r^+ . Operationally, bernoulli(x, y, p)generates a sample from the Bernoulli distribution with parameter p on the set $\{x, y\}$, whilst normal(x, y) generates a normal deviate with mean x and standard deviation y. We then postulate the metric axiom,

$$\frac{p, q \in [0, 1] \cap \mathbb{Q}}{\text{bernoulli}(x_1, x_2, p(*)) =_{|p-q|} \text{bernoulli}(x_1, x_2, q(*))}$$
(3)

We interpret the resulting linear metric λ -theory in the category Ban of Banach spaces and short operators, *i.e.* the semantics of [9, 20] without the order structure needed to interpret while loops. This is the usual representation of Markov chains/kernels as matrices/operators.

▶ **Theorem 29.** The category Ban is a Met-enriched autonomous category, and thus an instance of Definition 19.

In particular, Ban forms a model for the theory of our small probabilistic language via the following interpretation. We define $[[\text{Real}]] = \mathcal{M}\mathbb{R}$, the Banach space of finite Borel measures on \mathbb{R} equipped with the total variation norm, and similarly $[[\text{Real}^+]] = \mathcal{M}\mathbb{R}^+$ and $[[\text{unit}]] = \mathcal{M}[0, 1]$. We have $[[\mathbb{I}]] = \mathbb{R} \ni 1$, and for every $r \in \mathbb{Q}$ we put $[[r]] : \mathbb{R} \to \mathcal{M}\mathbb{R}, x \mapsto x\delta_r$, where δ_r is the Dirac delta over r; thus $[[r]](1) = \delta_r$. We define an analogous interpretation for the operation symbols r^+ and r^u . For $\mu, v \in \mathcal{M}\mathbb{R}$ we define $[[+]](\mu \otimes v) \triangleq +_*(\mu \otimes v)$ the pushforward under + of the product measure $\mu \otimes v$ (seen as an element of $\mathcal{M}\mathbb{R} \otimes \mathcal{M}\mathbb{R}$, see [9]). For $\mu, v, \xi \in \mathcal{M}\mathbb{R}$ we define $[[\texttt{bernoulli}]](\mu \otimes v \otimes \xi) \triangleq \text{bern}_*(\mu \otimes v \otimes \xi)$, the pushforward of the product measure $\mu \otimes v \otimes \xi$ under the Markov kernel bern : $\mathbb{R}^3 \to \mathbb{R}$, $(u, v, p) \mapsto p\delta_u + (1-p)\delta_v$, and similarly for [[normal]] (see [9] for the definition of pushforward by a Markov kernel).

This interpretation is sound (a proof is given in the Appendix) because the norm on $\mathcal{M}\mathbb{R}$ is the total variation norm, and the metric axiom (3) describes the total variation distance between the corresponding Bernoulli distributions. Consider now the following λ -terms (where we abbreviate the constants 0(*), 1(*), p(*), q(*) to 0, 1, p, q, respectively),

$$\begin{split} \texttt{walk1} &\triangleq \lambda x: \texttt{Real.bernoulli}(0, x + \texttt{normal}(0, 1), p) \\ \texttt{walk2} &\triangleq \lambda x: \texttt{Real.bernoulli}(0, x + \texttt{normal}(0, 1), q), \qquad p, q \in [0, 1] \cap \mathbb{Q}. \end{split}$$

As the names suggest, these two terms of type Real \rightarrow Real are denoted by random walks on \mathbb{R} . At each call, walk1 (resp. walk2) performs a jump drawn randomly from a standard normal distribution, or is forced to return to the origin with probability p (resp. q). These are non-standard random walks whose semantics are concretely given by complicated operators $\mathcal{M}\mathbb{R} \rightarrow \mathcal{M}\mathbb{R}$, but the simple quantitative equational system of Fig. 4 and the axiom (3) allow us to easily derive walk1 = $_{|p-q|}$ walk2 without having to compute the semantics of these terms. In other words, the soundness of (3) is enough to tightly bound the distance between two non-trivial random walks represented as higher-order terms in a probabilistic programming language. Furthermore, the tensor in the λ -calculus allows us to easily scale up this reasoning to random walks in higher dimensions such as walk1 \otimes walk2 on \mathbb{R}^2 .

5 Conclusions and future work

We introduced the notion of a \mathcal{V} -equation which generalises the well-established notions of equation, inequation [22, 2], and metric equation [29, 30]. We then presented a sound and complete \mathcal{V} -equational system for linear λ -calculus, illustrated with different examples of programs containing real-time and probabilistic behaviour.

Functorial connection to previous work. As a concluding note, let us introduce a simple yet instructive functorial connection between (1) the categorical semantics of linear λ -calculus with the \mathcal{V} -equational system, (2) the categorical semantics of linear λ -calculus with the equational system of Section 2, and (3) the algebraic semantics of the exponential free, multiplicative fragment of linear logic. First we need to recall some well-known facts. As detailed before, typical categorical models of linear λ -calculus and its equational system are locally small autonomous categories. The latter form a quasicategory Aut whose morphisms are autonomous functors. The usual algebraic models of the exponential free, multiplicative fragment of linear logic are the so-called lineales [10]. In a nutshell, a lineale is a poset (X, \leq) paired with a commutative, monoid operation $\otimes : X \times X \to X$ that satisfies certain conditions. Lineales are almost quantales: the only difference is that they do not require X to be cocomplete. The key idea in algebraic semantics is that the order \leq in the lineale encodes the logic's entailment relation. A functorial connection between autonomous categories and lineales (*i.e.* between (2) and (3)) is stated in [10] and is based on the following two

observations. First, (possibly large) lineales can be seen as *thin* autonomous categories, *i.e.* as elements of the enriched quasicategory $\{0, 1\}$ -Aut. Second, the inclusion $\{0, 1\}$ -Aut \hookrightarrow Aut has a left adjoint which *collapses all morphisms* of a given autonomous category C (intuitively, it eliminates the ability of C to differentiate different terms between two types). This provides an adjoint situation between (2) and (3). We can now expand this connection to our categorical semantics of linear λ -calculus and corresponding \mathcal{V} -equational system (*i.e.* (1)) in the following way. The forgetful functor \mathcal{V} -Cat \rightarrow Set has a left adjoint D: Set $\rightarrow \mathcal{V}$ -Cat which sends a set X to DX = (X, d),

$$d(x_1, x_2) = \begin{cases} k & \text{if } x_1 = x_2 \\ \bot & \text{otherwise} \end{cases}$$

This left adjoint is strong monoidal, specifically we have $D(X_1 \times X_2) = DX_1 \otimes DX_2$ and $\mathbb{I} = (1, (*, *) \mapsto k) = D1$. This gives rise to the functors,

$$(\mathcal{V}\text{-}\mathsf{Cat})\text{-}\mathsf{Aut} \xrightarrow{\hat{D}} \mathsf{Aut} \xrightarrow{c} \{0,1\}\text{-}\mathsf{Aut}$$

where \hat{D} equips the hom-sets of an autonomous category with the corresponding discrete \mathcal{V} -category and c collapses all morphisms of an autonomous category as described earlier. The right adjoint of \hat{D} forgets the \mathcal{V} -categorical structure between terms (*i.e.* morphisms) and the right adjoint of c is the inclusion functor mentioned earlier. Note that \hat{D} restricts to $(\mathcal{V}-\mathsf{Cat}_{\mathsf{sep}})$ -Aut and $(\mathcal{V}-\mathsf{Cat}_{\mathsf{sym},\mathsf{sep}})$ -Aut, and thus we obtain a functorial connection between the categorical semantics of linear λ -calculus with the \mathcal{V} -equational system (*i.e.* (1)), (2), and (3). In essence, the connection formalises the fact that our categorical models admit a richer structure over terms (*i.e.* morphisms) than the categorical models of linear λ -calculus and its classical equational system. The latter in turn permits the existence of different terms between two types as opposed to the algebraic semantics of the exponential free, multiplicative fragment of linear logic. The connection also shows that models for (2) and (3) can be mapped into models of our categorical semantics by equipping the respective hom-sets with a trivial, discrete structure.

Future work. Recall that linear λ -calculus is at the root of different ramifications of λ -calculus that relax resource-based conditions in different ways. Currently, we are studying analogous ramifications of linear λ -calculus in the \mathcal{V} -equational setting, particularly affine and Cartesian versions. We are also studying the possibility of adding an exponential modality in order to obtain a *mixed linear-non-linear* calculus [3]. We also started to explore different definitions of a morphism between $\mathcal{V}\lambda$ -theories and respective categories. This is the basis to establish a categorical equivalence between a (quasi)category of $\mathcal{V}\lambda$ -theories and a (quasi)category of \mathcal{V} -Cat_{sep}-enriched autonomous categories.

Next, our main examples of $\mathcal{V}\lambda$ -theories (see Section 4) used either the Boolean or the metric quantale. We would like to study linear $\mathcal{V}\lambda$ -theories whose underlying quantales are neither the Boolean nor the metric one, for example the ultrametric quantale which is (tacitly) used to interpret Nakano's guarded λ -calculus [5] and also to interpret a higher-order language for functional reactive programming [21]. Another interesting quantale is the Gödel one which is a basis for fuzzy logic [11] and whose \mathcal{V} -equations give rise to what we call fuzzy inequations.

Finally we plan to further explore the connections between our work and different results on metric universal algebra [29, 30, 35] and inequational universal algebra [22, 2, 35]. For example, an interesting connection is that the monad construction presented in [29] crucially relies on quotienting a pseudometric space into a metric space – this is a particular case of quotienting a \mathcal{V} -category into a separated \mathcal{V} -category (which we crucially use in our work).

— References

- Martín Abadi, Anindya Banerjee, Nevin Heintze, and Jon G Riecke. A core calculus of dependency. In Proceedings of the 26th ACM SIGPLAN-SIGACT symposium on Principles of programming languages, pages 147–160, 1999.
- 2 Jiří Adámek, Chase Ford, Stefan Milius, and Lutz Schröder. Finitary monads on the category of posets. arXiv preprint arXiv:2011.14796, 2020.
- 3 Nick Benton. A mixed linear and non-linear logic: Proofs, terms and models. In International Workshop on Computer Science Logic, pages 121–135. Springer, 1994.
- 4 Nick Benton, Gavin Bierman, Valeria de Paiva, and Martin Hyland. Term assignment for intuitionistic linear logic (preliminary report). Citeseer, 1992.
- 5 Lars Birkedal, Jan Schwinghammer, and Kristian Støvring. A metric model of lambda calculus with guarded recursion. In *FICS*, pages 19–25, 2010.
- 6 Francis Borceux. Handbook of categorical algebra, volume 2 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1994.
- 7 Raphaëlle Crubillé and Ugo Dal Lago. Metric reasoning about λ-terms: The affine case. In 2015 30th Annual ACM/IEEE Symposium on Logic in Computer Science, pages 633–644. IEEE, 2015.
- 8 Raphaëlle Crubillé and Ugo Dal Lago. Metric Reasoning About lambda-Terms: The General Case. In European Symposium on Programming, pages 341–367. Springer, 2017.
- 9 Fredrik Dahlqvist and Dexter Kozen. Semantics of higher-order probabilistic programs with conditioning. In Proc. 47th ACM SIGPLAN Symp. Principles of Programming Languages (POPL'20), pages 57:1–29, New Orleans, January 2020. ACM.
- 10 Valeria De Paiva. Lineales: algebraic models of linear logic from a categorical perspective. In Proceedings of LLC8, 1999.
- 11 Klaus Denecke, Marcel Erné, and Shelly L Wismath. Galois connections and applications, volume 565. Springer Science & Business Media, 2013.
- 12 Marco Gaboardi, Shin-ya Katsumata, Dominic Orchard, Flavien Breuvart, and Tarmo Uustalu. Combining effects and coeffects via grading. *ACM SIGPLAN Notices*, 51(9):476–489, 2016.
- 13 Francesco Gavazzo. Quantitative behavioural reasoning for higher-order effectful programs: Applicative distances. In Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, pages 452–461, 2018.
- 14 Gerhard Gierz, Karl Heinrich Hofmann, Klaus Keimel, Jimmie D. Lawson, Michael W. Mislove, and Dana S. Scott. *Continuous lattices and domains*, volume 93 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2003.
- 15 Jean-Yves Girard, Andre Scedrov, and Philip J Scott. Bounded linear logic: a modular approach to polynomial-time computability. *Theoretical computer science*, 97(1):1–66, 1992.
- 16 Jean Goubault-Larrecq. Non-Hausdorff Topology and Domain Theory—Selected Topics in Point-Set Topology, volume 22 of New Mathematical Monographs. Cambridge University Press, March 2013.
- 17 Dirk Hofmann and Pedro Nora. Hausdorff coalgebras. Applied Categorical Structures, 28(5):773– 806, 2020.
- 18 Shih-Han Hung, Kesha Hietala, Shaopeng Zhu, Mingsheng Ying, Michael Hicks, and Xiaodi Wu. Quantitative robustness analysis of quantum programs. Proceedings of the ACM on Programming Languages, 3(POPL):1–29, 2019.
- **19** Gregory Maxwell Kelly. *Basic concepts of enriched category theory*, volume 64. CUP Archive, 1982.
- 20 Dexter Kozen. Semantics of probabilistic programs. J. Comput. Syst. Sci., 22(3):328–350, June 1981. doi:10.1016/0022-0000(81)90036-2.
- 21 Neelakantan R Krishnaswami and Nick Benton. Ultrametric semantics of reactive programs. In 2011 IEEE 26th Annual Symposium on Logic in Computer Science, pages 257–266. IEEE, 2011.

- 22 Alexander Kurz and Jiří Velebil. Quasivarieties and varieties of ordered algebras: regularity and exactness. *Mathematical Structures in Computer Science*, 27(7):1153–1194, 2017.
- 23 Joachim Lambek and Philip J Scott. *Introduction to higher-order categorical logic*, volume 7. Cambridge University Press, 1988.
- 24 F William Lawvere. Metric spaces, generalized logic, and closed categories. Rendiconti del seminario matématico e fisico di Milano, 43(1):135–166, 1973.
- 25 Fred E. J. Linton. Some aspects of equational categories. In Proceedings of the Conference on Categorical Algebra, pages 84–94. Springer, 1966.
- 26 Saunders Mac Lane. Categories for the working mathematician, volume 5. springer, 1998.
- 27 Ian Mackie, Leopoldo Román, and Samson Abramsky. An internal language for autonomous categories. Applied Categorical Structures, 1(3):311–343, 1993.
- 28 Maria Emilia Maietti, Paola Maneggia, Valeria De Paiva, and Eike Ritter. Relating categorical semantics for intuitionistic linear logic. *Applied categorical structures*, 13(1):1–36, 2005.
- 29 Radu Mardare, Prakash Panangaden, and Gordon Plotkin. Quantitative algebraic reasoning. In Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, pages 700–709, 2016.
- 30 Radu Mardare, Prakash Panangaden, and Gordon Plotkin. On the axiomatizability of quantitative algebras. In 2017 32nd Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), pages 1–12. IEEE, 2017.
- 31 Dominic Orchard, Vilem-Benjamin Liepelt, and Harley Eades III. Quantitative program reasoning with graded modal types. *Proceedings of the ACM on Programming Languages*, 3(ICFP):1–30, 2019.
- 32 Jan Paseka and Jiří Rosický. Quantales. In Current research in operational quantum logic, pages 245–262. Springer, 2000.
- 33 Paolo Pistone. On Generalized Metric Spaces for the Simply Typed λ-Calculus. In Proceedings of the 36th Annual ACM/IEEE Symposium on Logic in Computer Science, 2021.
- 34 Jason Reed and Benjamin C Pierce. Distance makes the types grow stronger: A calculus for differential privacy. In Proceedings of the 15th ACM SIGPLAN international conference on Functional programming, pages 157–168, 2010.
- 35 Jiří Rosický. Metric monads. arXiv preprint arXiv:2012.14641, 2020.
- 36 Raymond A Ryan. Introduction to tensor products of Banach spaces. Springer Science & Business Media, 2013.
- 37 Michael Shulman. A practical type theory for symmetric monoidal categories. *arXiv preprint arXiv:1911.00818*, 2019.
- 38 Isar Stubbe. An introduction to quantaloid-enriched categories. Fuzzy Sets and Systems, 256:95–116, 2014.
- 39 Jiří Velebil, Alexander Kurz, and Adriana Balan. Extending set functors to generalised metric spaces. Logical Methods in Computer Science, 15, 2019.

A Linear $\mathcal{V}\lambda$ -theories and linear quantales

We briefly study linear $\mathcal{V}\lambda$ -theories where \mathcal{V} is a quantale with a *linear order*. The latter condition is respected by the Boolean and (ultra)metric quantales (mentioned in the main text). Recall that metric universal algebra [29, 30] tacitly uses the metric quantale.

Theorem 30. Assume that the underlying order of \mathcal{V} is linear and consider a (symmetric) linear $\mathcal{V}\lambda$ -theory. Substituting the rule below on the left by the one below on the right does not change the theory.

$$\frac{\forall i \le n. \ v =_{q_i} w}{v =_{\lor q_i} w} \qquad \qquad \overline{v =_{\bot} w}$$

Proof. Clearly, the rule on the left subsumes the one on the right by choosing n = 0. So we only need to show the inverse direction under the assumption that \mathcal{V} is linear. Thus, assume that $\forall i \leq n$. $v =_{q_i} w$. We proceed by case distinction. If n = 0 then we need to show that $v =_{\perp} w$ which is given already by the rule on the right. Suppose now that n > 0. Then since the order of \mathcal{V} is linear the value $\forall q_i$ must already be one of the values q_i and $v =_{q_i} w$ is already part of the theory. In other words, in case of n > 0 the rule on the left is redundant.

The above result is in accordance with metric universal algebra [29, 30] which also does not include rule (join). Interestingly, however, we still have $v =_{\perp} w$ for all λ -terms v and wand such a rule is not present in [29, 30]. This is explained by the fact that metric equations in [29, 30] are labelled *only* by non-negative rational numbers whilst we also permit infinity to be a label (in our case, labels are given by a basis B which for the metric case corresponds to the *extended* non-negative rational numbers). All remaining rules of our \mathcal{V} -equational system instantiated to the metric case find a counterpart in the metric equational system presented in [29, 30].

Next, note that if the quantale \mathcal{V} is finite then for all $q \in \mathcal{V}$ we have $q \ll q$ which means that rule (arch) is no longer necessary. This observation is applicable to the Boolean quantale.

B Lemmata and omitted proofs

Proof sketch of Lemma 1. The proof follows by induction over the structure of λ -terms. Here we only consider the case $f(v_1, \ldots, v_n)$, because the other cases follow analogously.

Suppose that $E \triangleright f(v_1, \ldots, v_n) : \mathbb{A}$. Then according to the typing system it is necessarily the case that the previous derivations were $\Gamma_i \triangleright v_i : \mathbb{A}$ for all $i \leq n$ with $E \in \mathrm{Sf}(\Gamma_1; \ldots; \Gamma_n)$ for some family of contexts $(\Gamma_i)_{i \leq n}$. The only room for choice is therefore in choosing the contexts Γ_i . We will show that even this choice is unique. Consider two families $(\Gamma_i)_{i \leq n}$ and $(\Gamma'_i)_{i \leq n}$ such that $\Gamma_i \triangleright v_i : \mathbb{A}$ and $\Gamma'_i \triangleright v_i : \mathbb{A}$ for all $i \leq n$, and moreover $E \in \mathrm{Sf}(\Gamma_1; \ldots; \Gamma_n)$ and $E \in \mathrm{Sf}(\Gamma'_1; \ldots; \Gamma'_n)$. Since $\Gamma_i \triangleright v_i : \mathbb{A}_i$ and $\Gamma'_i \triangleright v_i : \mathbb{A}_i$ we deduce (by linearity) that Γ_i is a permutation of Γ'_i . Consequently, since $E \in \mathrm{Sf}(\Gamma_1, \ldots, \Gamma_n)$, $E \in \mathrm{Sf}(\Gamma'_1, \ldots, \Gamma'_n)$ and E (by the definition of a shuffle) cannot change the relative order of the elements in Γ_i and Γ'_i for all $i \leq n$, it must be the case that $\Gamma_i = \Gamma'_i$ for all $i \leq n$. In other words, the choice of $(\Gamma_i)_{i \leq n}$ is fixed a priori. The proof now follows by applying the induction hypothesis to each v_i .

Proof sketch of Lemma 2. We focus first on the exchange rule. The proof follows by induction over the structure of derivations. Here we only consider the case $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \triangleright f(v_1, \ldots, v_n) : \mathbb{C}$, the other cases follow analogously.

Suppose that $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \triangleright f(v_1, \dots, v_n) : \mathbb{C}$ with $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \in \mathrm{Sf}(\Gamma_1; \dots; \Gamma_n)$. We proceed by case distinction: assume first that both $x : \mathbb{A}$ and $y : \mathbb{B}$ are in some Γ_i , with $i \leq n$. We can thus decompose Γ_i into $\Gamma_i^1, x : \mathbb{A}, y : \mathbb{B}, \Gamma_i^2$. Then we apply the induction hypothesis on $\Gamma_i \triangleright v_i : \mathbb{A}_i$ and proceed by observing that if $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \in \mathrm{Sf}(\Gamma_1; \dots; (\Gamma_i^1, x : \mathbb{A}, y : \mathbb{B}, \Gamma_i^2); \dots; \Gamma_n)$ then it is also the case that $\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \in \mathrm{Sf}(\Gamma_1; \dots; (\Gamma_i^1, y : \mathbb{B}, x : \mathbb{A}, \Gamma_i^2); \dots; \Gamma_n)$. Assume now that $x : \mathbb{A}$ is in some Γ_i and $y : \mathbb{B}$ is in some Γ_j with $i \neq j$. Then since $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \in \mathrm{Sf}(\Gamma_1; \dots; \Gamma_n)$ it must be the case that $\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \in \mathrm{Sf}(\Gamma_1; \dots; \Gamma_n)$ so we only need to apply rule (**ax**).

Let us now focus on the substitution rule. The proof follows by induction over the structure of derivations and also by the exchange rule that was just proved. We exemplify this with rule (\mathbf{ax}) . The other cases follow analogously.

Suppose that $\Gamma, x : \mathbb{A} \rhd f(v_1, \ldots, v_n) : \mathbb{B}$. Then for all $i \leq n$ we have $\Gamma_i \rhd v_i : \mathbb{A}_i$ and $\Gamma, x : \mathbb{A} \in \mathrm{Sf}(\Gamma_1; \ldots; \Gamma_n)$. By linearity and by the definition of a shuffle there exists exactly one Γ_i that can be decomposed into $\Gamma_i = \Gamma'_i, x : \mathbb{A}$. We then use the induction hypothesis to obtain $\Gamma'_i, \Delta \rhd v_i[w/x] : \mathbb{A}_i$. Now observe that if $\Gamma, x : \mathbb{A} \in \mathrm{Sf}(\Gamma_1; \ldots; (\Gamma'_i, x : \mathbb{A}); \ldots; \Gamma_n)$ then $\Gamma, \Delta \in \mathrm{Sf}(\Gamma_1; \ldots; (\Gamma'_i, \Delta); \ldots; \Gamma_n)$. We use this last observation to build $\Gamma, \Delta \rhd f(v_1, \ldots, v_i[w/x], \ldots, v_n) = f(v_1, \ldots, v_n)[w/x] : \mathbb{B}$.

In order to keep calculations in the following proofs legible we will sometimes abbreviate a denotation $\llbracket \Gamma \rhd v : \mathbb{A} \rrbracket$ to $\llbracket \Gamma \rhd v \rrbracket$ or even just $\llbracket v \rrbracket$.

▶ Lemma 31 (Exchange and Substitution). Consider judgements $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \triangleright v : \mathbb{C},$ $\Gamma, x : \mathbb{A} \triangleright v : \mathbb{B}, and \Delta \triangleright w : \mathbb{A}.$ Then the following equations hold in every autonomous category C:

$$\begin{split} \llbracket \Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \rhd v : \mathbb{C} \rrbracket = \llbracket \Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta \rhd v : \mathbb{C} \rrbracket \cdot \mathsf{exch}_{\Gamma,\underline{\mathbb{A}},\mathbb{B},\Delta} \\ \llbracket \Gamma, \Delta \rhd v[w/x] : \mathbb{B} \rrbracket = \llbracket \Gamma, x : \mathbb{A} \rhd v : \mathbb{B} \rrbracket \cdot \mathsf{jn}_{\Gamma:\mathbb{A}} \cdot (\mathsf{id} \otimes \llbracket \Delta \rhd w : \mathbb{A} \rrbracket) \cdot \mathsf{sp}_{\Gamma:\Delta} \end{split}$$

Proof sketch. For both cases the proof follows by induction over the structure of derivations. Here we only consider rule (\mathbf{ax}) , because the other ones follow analogously. In many of the calculations below we will tacitly perform simple diagram chases that take advantage of naturality, functoriality, and the coherence theorem of symmetric monoidal categories.

We start with the exchange property. Suppose that $\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \triangleright f(v_1, \ldots, v_n) : \mathbb{C}$. We proceed by case distinction: first, consider the case in which $x : \mathbb{A} \in \Gamma_i$ and $y : \mathbb{B} \in \Gamma_j$ with $i \neq j$. The proof then follows directly by observing that the two corresponding shuffling morphisms $\mathsf{sh}_{\Gamma,\mathbb{A},\mathbb{B},\Delta} : [\![\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta]\!] \to [\![\Gamma_1, \ldots, \Gamma_n]\!]$ and $\mathsf{sh}_{\Gamma,\mathbb{B},\mathbb{A},\Delta} : [\![\Gamma, y : \mathbb{B}, x : \mathbb{A}, \Delta]\!] \to$ $[\![\Gamma_1, \ldots, \Gamma_n]\!]$ satisfy the equation $\mathsf{sh}_{\Gamma,\mathbb{B},\mathbb{A},\Delta} \cdot \mathsf{exch}_{\Gamma,\mathbb{A},\mathbb{B},\Delta} = \mathsf{sh}_{\Gamma,\mathbb{A},\mathbb{B},\Delta}$. Consider now the case in which $x : \mathbb{A} \in \Gamma_i$ and $y : \mathbb{B} \in \Gamma_i$ for some $i \leq n$. We then calculate:

$$\begin{split} & [\![\Gamma, x : \mathbb{A}, y : \mathbb{B}, \Delta \rhd f(v_1, \dots, v_n) : \mathbb{C}]\!] = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes [\![v_i]\!] \otimes \dots \otimes [\![v_n]\!]) \cdot \mathsf{sp}_{\Gamma_1; \dots; \Gamma_n} \cdot \mathsf{sh}_{\Gamma, \mathbb{A}, \mathbb{B}, \Delta} \\ & = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes ([\![v_i]\!] \cdot \mathsf{exch}_{\Gamma_i^1, \underline{\mathbb{A}}, \underline{\mathbb{B}}, \Gamma_i^2}) \otimes \dots \otimes [\![v_n]\!]) \cdot \mathsf{sp}_{\Gamma_1; \dots; \Gamma_n} \cdot \mathsf{sh}_{\Gamma, \mathbb{A}, \mathbb{B}, \Delta} \\ & = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes [\![v_n]\!]) \cdot (\mathsf{id} \otimes \dots \otimes \mathsf{exch}_{\Gamma_i^1, \underline{\mathbb{A}}, \underline{\mathbb{B}}, \Gamma_i^2} \otimes \dots \otimes \mathsf{id}) \cdot \mathsf{sp}_{\Gamma_1; \dots; \Gamma_n} \cdot \mathsf{sh}_{\Gamma, \mathbb{A}, \mathbb{B}, \Delta} \\ & = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes [\![v_n]\!]) \cdot \mathsf{sp}_{\Gamma_1'; \dots; \Gamma_n'} \cdot \mathsf{exch}_{\Gamma_1, \dots, \Gamma_i^1, \underline{\mathbb{A}}, \underline{\mathbb{B}}, \Gamma_i^2, \dots, \Gamma_n} \cdot \mathsf{sh}_{\Gamma, \mathbb{A}, \mathbb{B}, \Delta} \\ & = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes [\![v_n]\!]) \cdot \mathsf{sp}_{\Gamma_1'; \dots; \Gamma_n'} \cdot \mathsf{sh}_{\Gamma, \mathbb{B}, \mathbb{A}, \Delta} \cdot \mathsf{exch}_{\Gamma, \underline{\mathbb{A}}, \underline{\mathbb{B}}, \Delta} \\ & = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes [\![v_n]\!]) \cdot \mathsf{sp}_{\Gamma_1'; \dots; \Gamma_n'} \cdot \mathsf{sh}_{\Gamma, \mathbb{B}, \mathbb{A}, \Delta} \cdot \mathsf{exch}_{\Gamma, \underline{\mathbb{A}}, \mathbb{B}, \Delta} \\ & = [\![f]\!] \cdot ([\![v_1]\!] \otimes \dots \otimes [\![v_n]\!]) \cdot \mathsf{sp}_{\Gamma_1'; \dots; \Gamma_n'} \cdot \mathsf{sh}_{\Gamma, \mathbb{B}, \mathbb{A}, \Delta} \cdot \mathsf{exch}_{\Gamma, \underline{\mathbb{A}}, \mathbb{B}, \Delta} \end{split}$$

Let us now focus on proving the substitution lemma for rule (**ax**):

$$\begin{split} & \left[\!\!\left[\Gamma,\Delta \rhd f(v_1,\ldots,v_n)[w/x] : \mathbb{B}\right]\!\!\right] = \left[\!\!\left[\Gamma,\Delta \rhd f(v_1,\ldots,v_i[w/x],\ldots,v_n)\right]\!\!\right] \\ &= \left[\!\!\left[f\right]\!\!\right] \cdot \left(\!\left[\!\!\left[v_1\right]\!\right] \otimes \cdots \otimes \left[\!\!\left[v_i[w/x]\right]\!\right] \otimes \cdots \otimes \left[\!\!\left[v_n\right]\!\right]\!\right) \cdot \mathsf{sp}_{\Gamma_1;\ldots;\Gamma_n} \cdot \mathsf{sh}_{\Gamma,\Delta} \\ &= \left[\!\!\left[f\right]\!\!\right] \cdot \left(\!\left[\!\!\left[v_1\right]\!\right] \otimes \cdots \otimes \left(\!\left[\!\!\left[v_i\right]\!\right] \cdot \mathsf{jn}_{\Gamma'_i;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!\left[w\right]\!\right]\!\right) \cdot \mathsf{sp}_{\Gamma'_i;\Delta}\right) \otimes \cdots \otimes \left[\!\!\left[v_n\right]\!\right] \right) \cdot \mathsf{sp}_{\Gamma_1;\ldots;\Gamma_n} \cdot \mathsf{sh}_{\Gamma,\Delta} \\ &= \left[\!\!\left[f\right]\!\right] \cdot \left(\!\left[\!\!\left[v_1\right]\!\right] \otimes \cdots \otimes \left[\!\!\left[v_n\right]\!\right]\right) \cdot \left(\mathsf{id} \otimes \cdots \otimes \left(\!\!\left[\mathsf{jn}_{\Gamma'_i;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!\left[w\right]\!\right]\!\right) \cdot \mathsf{sp}_{\Gamma'_i;\Delta}\right) \otimes \cdots \otimes \mathsf{id}\right) \cdot \mathsf{sp}_{\Gamma_1;\ldots;\Gamma_n} \cdot \mathsf{sh}_{\Gamma,\Delta} \\ &= \left[\!\!\left[f\right]\!\right] \cdot \left(\!\left[\!\!\left[v_1\right]\!\right] \otimes \cdots \otimes \left[\!\!\left[v_n\right]\!\right]\right) \cdot \mathsf{sp}_{\Gamma_1;\ldots;\Gamma'_i,\mathbb{A};\ldots,\Gamma_n} \cdot \mathsf{sh}_{\Gamma,\mathbb{A}} \cdot \mathsf{jn}_{\Gamma;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!\left[w\right]\!\right]\right) \cdot \mathsf{sp}_{\Gamma;\Delta} \\ &= \left[\!\!\left[\Gamma,x : \mathbb{A} \rhd f(v_1,\ldots,v_n)\!\right]\!\right] \cdot \mathsf{jn}_{\Gamma;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!w\!\right]\!\right]) \cdot \mathsf{sp}_{\Gamma;\Delta} \end{split}$$

Proof sketch of Theorem 3. The proof follows by an appeal to Lemma 31, the coherence theorem for symmetric monoidal categories, and naturality. We exemplify this with the commuting conversions.

$$\begin{split} & \left[\!\left[\Gamma, \Delta, E \triangleright u[v \text{ to } * . w/x] : \mathbb{B}\right]\!\right] = \left[\!\left[u\right]\!\right] \cdot \mathsf{jn}_{\Gamma;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\left[v \text{ to } * . w\right]\!\right]) \cdot \mathsf{sp}_{\Gamma;\Delta,E} \\ & = \left[\!\left[u\right]\!\right] \cdot \mathsf{jn}_{\Gamma;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\left[w\right]\!\right]) \cdot \lambda \cdot \left(\left[\!\left[v\right]\!\right] \otimes \mathsf{id}\right) \cdot \mathsf{sp}_{\Delta;E}\right)\right) \cdot \mathsf{sp}_{\Gamma;\Delta,E} \\ & = \left[\!\left[u\right]\!\right] \cdot \mathsf{jn}_{\Gamma;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\left[w\right]\!\right]) \cdot (\mathsf{id} \otimes (\lambda \cdot \left(\left[\!\left[v\right]\!\right] \otimes \mathsf{id}\right) \cdot \mathsf{sp}_{\Delta;E})\right) \cdot \mathsf{sp}_{\Gamma;\Delta,E} \\ & = \left[\!\left[u\right]\!\right] \cdot \mathsf{jn}_{\Gamma;\mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\left[w\right]\!\right]\right) \cdot \mathsf{sp}_{\Gamma;E} \cdot \mathsf{jn}_{\Gamma;E} \cdot (\mathsf{id} \otimes (\lambda \cdot \left(\left[\!\left[v\right]\!\right] \otimes \mathsf{id}\right) \cdot \mathsf{sp}_{\Delta;E})\right) \cdot \mathsf{sp}_{\Gamma;\Delta,E} \\ & = \left[\!\left[u[w/x]\right]\!\right] \cdot \mathsf{jn}_{\Gamma;E} \cdot (\mathsf{id} \otimes (\lambda \cdot \left(\left[\!\left[v\right]\!\right] \otimes \mathsf{id}\right) \cdot \mathsf{sp}_{\Delta;E})\right) \cdot \mathsf{sp}_{\Gamma;\Delta,E} \\ & = \left[\!\left[u[w/x]\right]\!\right] \cdot \lambda \cdot \left(\left[\!\left[v\right]\!\right] \otimes \mathsf{id}\right) \cdot \mathsf{sp}_{\Delta;\Gamma,E} \cdot \mathsf{sh}_{\Gamma,\Delta,E} \\ & = \left[\!\left[\Gamma, \Delta, E \triangleright v \text{ to } * . u[w/x]\right] : \mathbb{B}\!\right] \end{split}$$

The one but last step amounts to a diagram chase that recurs to naturality and the coherence theorem of symmetric monoidal categories.

$$\begin{split} & \left[\!\!\left[\Gamma, \Delta, E \rhd u[\operatorname{pm} v \text{ to } x \otimes y. \; w/z] : \mathbb{B}\!\right] = \left[\!\!\left[\Gamma, z : A \rhd u\!\right] \cdot \mathsf{jn}_{\Gamma; \mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!\left[\operatorname{pm} v \text{ to } x \otimes y. \; w\!\right]\!\right] \cdot \mathsf{sp}_{\Gamma; \Delta, E} \\ & = \left[\!\!\left[u\!\right]\!\right] \cdot \mathsf{jn}_{\Gamma; \mathbb{A}} \cdot (\mathsf{id} \otimes \left(\!\left[\!\!\left[w\!\right]\!\right] \cdot \mathsf{jn}_{E; \mathbb{C}; \mathbb{D}} \cdot \alpha \cdot \mathsf{sw} \cdot \left(\!\left[\!\!\left[v\!\right]\!\right] \otimes \mathsf{id}\!\right] \cdot \mathsf{sp}_{\Delta, E})\right) \cdot \mathsf{sp}_{\Gamma; \Delta, E} \\ & = \left[\!\!\left[u\!\right]\!\right] \cdot \mathsf{jn}_{\Gamma; \mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!\left[w\!\right]\!\right]) \cdot (\mathsf{id} \otimes (\mathsf{jn}_{E; \mathbb{C}; \mathbb{D}} \cdot \alpha \cdot \mathsf{sw} \cdot \left(\!\left[\!\!\left[v\!\right]\!\right] \otimes \mathsf{id}\!\right] \cdot \mathsf{sp}_{\Delta; E})\right) \cdot \mathsf{sp}_{\Gamma; \Delta, E} \\ & = \left[\!\!\left[u\!\right]\!\right] \cdot \mathsf{jn}_{\Gamma; \mathbb{A}} \cdot (\mathsf{id} \otimes \left[\!\!\left[w\!\right]\!\right]) \cdot \mathsf{sp}_{\Gamma; E, \mathbb{C}, \mathbb{D}} \cdot \mathsf{jn}_{\Gamma; E, \mathbb{C}, \mathbb{D}} \cdot (\mathsf{id} \otimes (\mathsf{jn}_{E; \mathbb{C}; \mathbb{D}} \cdot \alpha \cdot \mathsf{sw} \cdot \left(\!\left[\!\!\left[v\!\right]\!\right] \otimes \mathsf{id}\!\right] \cdot \mathsf{sp}_{\Delta; E})\right) \cdot \mathsf{sp}_{\Gamma; \Delta, E} \\ & = \left[\!\!\left[u\!\left[w/z\!\right]\!\right] \cdot \mathsf{jn}_{\Gamma; E, \mathbb{C}, \mathbb{D}} \cdot (\mathsf{id} \otimes (\mathsf{jn}_{E; \mathbb{C}; \mathbb{D}} \cdot \alpha \cdot \mathsf{sw} \cdot (\left[\!\!\left[v\!\right]\!\right] \otimes \mathsf{id}\!\right) \cdot \mathsf{sp}_{\Delta; E})\right) \cdot \mathsf{sp}_{\Gamma; \Delta, E} \\ & = \left[\!\!\left[u\!\left[w/z\!\right]\!\right] \cdot \mathsf{jn}_{\Gamma, E; \mathbb{C}; \mathbb{D}} \cdot \alpha \cdot \mathsf{sw} \cdot (\left[\!\!\left[v\!\right]\!\right] \otimes \mathsf{id}\!\right) \cdot \mathsf{sp}_{\Delta; \Gamma, E} \cdot \mathsf{sh}_{\Gamma, \Delta, E} \\ & = \left[\!\!\left[\Gamma, \Delta, E \triangleright \mathsf{pm} v \text{ to } x \otimes y. \; u\!\left[\!w/z\!\right]\!\right] : \mathbb{B}\!\right] \end{split}$$

The one but last step amounts to a diagram chase that recurs to naturality and the coherence theorem of symmetric monoidal categories.

Proof of Theorem 17. The proof follows by showing that the closed monoidal structure of \mathcal{V} -Cat preserves symmetry and separation. It is immediate for symmetry. For separation, note that since \mathcal{V} is integral the inequation $x \otimes y \leq x$ holds for all $x, y \in \mathcal{V}$. It follows that the monoidal structure preserves separation. The fact that the closed structure also preserves separation uses the implication $x \leq \bigwedge A \Rightarrow \forall a \in A. x \leq a$ for all $x \in X, A \subseteq X$.

Proof of Theorem 25. Let us denote by $\operatorname{Lang}^{\lambda}(\mathsf{C})$ the *linear* λ -theory generated from C . According to Theorem 8, the category $\operatorname{Syn}(\operatorname{Lang}^{\lambda}(\mathsf{C}))$ (*i.e.* the syntactic category generated from $\operatorname{Lang}^{\lambda}(\mathsf{C})$) is locally small whenever C is locally small. Then consider two types \mathbb{A} and

 \mathbb{B} . We will prove our claim by taking advantage of the axiom of replacement in ZF set-theory, specifically by presenting a *surjective* map,

$$\operatorname{Syn}(\operatorname{Lang}^{\lambda}(\mathsf{C}))(\mathbb{A},\mathbb{B})\longrightarrow \operatorname{Syn}(\operatorname{Lang}(\mathsf{C}))(\mathbb{A},\mathbb{B})$$

The crucial observation is that if v = w in $\text{Lang}^{\lambda}(\mathsf{C})$ then $v =_{\top} w$ and $w =_{\top} v$ in $\text{Lang}(\mathsf{C})$. This is obtained by the definition of a model, the definition of a \mathcal{V} -category, and the definition of $\text{Lang}(\mathsf{C})$. This observation allows to establish the surjective map that sends [v] to [v], *i.e.* it sends the equivalence class of v as a λ -term in $\text{Lang}^{\lambda}(\mathsf{C})$ into the equivalence class of v as a λ -term in $\text{Lang}^{\lambda}(\mathsf{C})$.

Proof of Theorem 29. The autonomous structure of Ban is well-known [9, 20], so let us focus on showing that it is a Met-enriched autonomous category. The enrichment is simply given by distance function induced by the operator norm, thus if $S, T \in Ban(X, Y)$,

$$d(S,T) = \bigvee \{ \| (S-T)(x) \| \mid \|x\| \le 1 \}$$

Composition is a short map.

Let $T, T' \in Ban(X, Y)$ and $S, S' \in Ban(Y, Z)$, we compute:

$$\begin{split} d(ST, S'T') &\triangleq \bigvee \{ \|ST(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &= \bigvee \{ \|ST(x) - ST'(x) + ST'(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &\le \bigvee \{ \|ST(x) - ST'(x)\| + \|ST'(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &= \bigvee \{ \|ST(x) - ST'(x)\| \mid \|x\| \le 1 \} + \bigvee \{ \|ST'(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &= \bigvee \{ \|S(T - T')(x)\| \mid \|x\| \le 1 \} + \bigvee \{ \|ST'(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &\le \bigvee \{ \|T(x) - T'(x)\| \mid \|x\| \le 1 \} + \bigvee \{ \|ST'(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &\le \bigvee \{ \|T(x) - T'(x)\| \mid \|x\| \le 1 \} + \bigvee \{ \|ST'(x) - S'T'(x)\| \mid \|x\| \le 1 \} \\ &\le \bigcup \{ \|T(x) - T'(x)\| \mid \|x\| \le 1 \} + \bigvee \{ \|S(y) - S'(y)\| \mid \|y\| \le 1 \} \\ &\triangleq d(T, T') + d(S, S') \\ &\triangleq d((T, S), (T', S')) \end{split}$$

where (\star) follows from the fact that,

$$\|S(T - T')(x)\| = \|T(x) - T'(x)\| \left\| S\left(\frac{(T - T')(x)}{\|T(x) - T'(x)\|}\right) \right\| \le \|T(x) - T'(x)\|$$

by linearity of S and by the fact that $||S|| \leq 1$. This shows that Ban is Met-enriched. We now turn to the first clause of Definition 19.

The monoidal operation is an enriched bi-functor.

Note first that if $S \in \mathsf{Ban}(X, Y)$ and $T, T' \in \mathsf{Ban}(X', Y')$ then,

 $S\widehat{\otimes}_{\pi}T - S\widehat{\otimes}_{\pi}T' = S\widehat{\otimes}_{\pi}(T - T')$

Indeed, since $S \widehat{\otimes}_{\pi} T$ is the unique linear operator such that $S \widehat{\otimes}_{\pi} T(x \otimes x') = S(x) \otimes T(x')$, we can reason pointwise and get,

$$(S\widehat{\otimes}_{\pi}T - S\widehat{\otimes}_{\pi}T')(x \otimes x') = (S\widehat{\otimes}_{\pi}T)(x \otimes x') - (S\widehat{\otimes}_{\pi}T')(x \otimes x')$$
$$= S(x) \otimes T(x') - S(x) \otimes T'(x')$$
$$= S(x) \otimes (T(x') - T'(x'))$$
$$= (S\widehat{\otimes}_{\pi}(T - T'))(x \otimes x')$$

where the penultimate step follows from the basic definition of the tensor product of vector spaces. Now we can show that for any Banach spaces X, X', Y, Y' the projective tensor map,

$$\mathsf{Ban}(X,Y) \otimes \mathsf{Ban}(X',Y') \to \mathsf{Ban}(X \widehat{\otimes}_{\pi} X',Y \widehat{\otimes}_{\pi} Y'), (S,T) \mapsto S \widehat{\otimes}_{\pi} T$$

where \otimes once again denotes the monoidal operation in Met, is short. We simply compute,

$$d(S\widehat{\otimes}_{\pi}T, S'\widehat{\otimes}_{\pi}T') \triangleq \|S\widehat{\otimes}_{\pi}T - S'\widehat{\otimes}_{\pi}T'\|$$

$$= \|S\widehat{\otimes}_{\pi}T - S\widehat{\otimes}_{\pi}T' + S\widehat{\otimes}_{\pi}T' - S'\widehat{\otimes}_{\pi}T'\|$$

$$\leq \|S\widehat{\otimes}_{\pi}T - S\widehat{\otimes}_{\pi}T'\| + \|S\widehat{\otimes}_{\pi}T' - S'\widehat{\otimes}_{\pi}T'\|$$

$$= \|S\widehat{\otimes}_{\pi}(T - T')\| + \|(S - S')\widehat{\otimes}_{\pi}T'\|$$

$$= \|S\|\|T - T'\| + \|S - S'\|\|T'\|$$

$$\leq \|T - T'\| + \|S - S'\| \triangleq d((S, T), (S, T'))$$

where the last step uses the fact that $||S||, ||T|| \leq 1$ and the penultimate step uses the basic fact that $||S \widehat{\otimes}_{\pi} T|| = ||S|| ||T||$ (see [36, §2.1]). Finally, we show the second clause of Definition 19.

The adjunction $-\widehat{\otimes}_{\pi}Y \vdash Y \multimap -$ is a Met-adjunction. The fact that the maps,

$$\mathsf{Ban}(X, X') \to \mathsf{Ban}(X \widehat{\otimes}_{\pi} Y, X \widehat{\otimes}_{\pi} Y), \ f \mapsto f \widehat{\otimes}_{\pi} \operatorname{id}_{Y}$$

are short follows by re-writing them as,

$$\mathsf{Ban}(X, X') \simeq \mathsf{Ban}(X, X') \otimes 1 \longrightarrow \mathsf{Ban}(X, X') \otimes \mathsf{Ban}(Y, Y) \longrightarrow \mathsf{Ban}(X \widehat{\otimes}_{\pi} X', Y \widehat{\otimes}_{\pi} Y)$$

and the fact that the monoidal operation of Ban is an enriched bi-functor. Similarly, the map,

$$\mathsf{Ban}(X,X')\to\mathsf{Ban}(Y\multimap X,Y\multimap X'),\ S\mapsto (T\mapsto ST)$$

is short. This is a consequence of the following fact. Consider two operators $S, S' \in Ban(X, X')$. For all bounded operators $T \in Y \multimap X$ with $||T|| \leq 1$ we have $d(S, S') \triangleq ||S - S'|| \geq ||ST - S'T||$, which provides,

$$d(S,S') \ge \bigvee \{ \|ST - S'T\| \mid \|T\| \le 1 \} \triangleq d(T \mapsto ST, T \mapsto S'T)$$

Finally, we need to show that the adjunction $-\widehat{\otimes}_{\pi}Y \vdash Y \multimap -$ defines an isometry,

$$\mathsf{Ban}(X\widehat{\otimes}_{\pi}Y,Z)\simeq\mathsf{Ban}(X,Y\multimap Z).$$

Indeed, the bijection from left to right is defined by,

$$T \mapsto (Y \multimap -)(T) \cdot \eta_Y$$

where η is the unit of the adjunction. Since $Y \multimap -$ is Met-enriched, the assignment $T \mapsto (Y \multimap -)(T)$ is short, and composition by η_Y is short. Thus the invertible map $\operatorname{Ban}(X \widehat{\otimes}_{\pi} Y, Z) \to \operatorname{Ban}(X, Y \multimap Z)$ is short. By a similar argument using the co-unit of the adjunction and the fact that $-\widehat{\otimes}_{\pi} Y$ is Met-enriched we get that the invertible map $\operatorname{Ban}(X, Y \multimap Z) \to \operatorname{Ban}(X \widehat{\otimes}_{\pi} Y, Z)$ is also short. It follows that both maps must be invertible isometries.

Proof that the axiom (3) is sound.

The total variation distance between [[bernoulli(x, y, p)]] and [[bernoulli(x, y, q)]] with $p, q \in [0, 1]$ is given by:

$$\begin{split} &\bigvee_{A} \left| \iint p\delta_{u}(A) + (1-p)\delta_{v}(A) \ d\llbracket x \rrbracket (du)d\llbracket y \rrbracket (dv) - \iint q\delta_{u}(A) + (1-q)\delta_{v}(A) \ d\llbracket x \rrbracket (du)d\llbracket y \rrbracket (dv) \right| \\ &= \bigvee_{A} \left| \iint (p-q)\delta_{u}(A) + ((1-p) - (1-q))\delta_{v}(A) \ d\llbracket x \rrbracket (du)d\llbracket y \rrbracket (dv) \right| \\ &= \bigvee_{A} |(p-q)\llbracket x \rrbracket (A) + (q-p)\llbracket y \rrbracket (A)| \\ &= \bigvee_{A} |(p-q)(\llbracket x \rrbracket (A) - \llbracket y \rrbracket (A))| \\ &\leq |p-q| \end{split}$$